

Cordination Compound of Dimethyltin(IV) with N,N,N',N'-Tetraethylethylenediamine: Speciation and Theoretical approach

Safaa Said Hassan¹, Mohamed Mohamed Shoukry^{*1,2}, Abdel Aziz Qasem Jbarah³

¹Department of Chemistry, Faculty of Science, Cairo University, Giza, Egypt.

²Department of Chemistry, Faculty of Science, Islamic University, Al-Madinah, Saudi Arabia.

³Chemistry Department, College of Science, Al-Hussein Bin Talal University, Ma'an, Jordan.

***Corresponding author:** Mohamed Mohamed Shoukry, email: shoukrymm@hotmail.com

Received August 6th, 2019; Accepted January 21th, 2020.

DOI: <http://dx.doi.org/10.29356/jmcs.v64i2.1079>

Abstract. The formation equilibria of the dimethyltin(IV) complexes with of N,N,N',N'-tetraethylethylenediamine (Et₄en) in solution were investigated. The stoichiometry and stability constants of the complexes formed in solution phase were determined at different temperatures (15 °C – 35 °C) and in solutions of dioxane-water mixtures of different compositions (15% - 62.5%). The accepted model is composed of the 110, 111, 11-1 and 11-2 species. The thermodynamic parameters ΔH° and ΔS° associated with the protonation of N,N,N',N'-tetraethylethylenediamine (Et₄en) and its complex formation with the dimethyltin(IV) species were determined. The complex formation reaction is exothermic. The equilibrium constant for the displacement of N,N,N',N'-tetraethylethylenediamine coordinated to dimethyltin(IV) by some selected DNA constituents was calculated. The K_{eq} values clearly indicate the ability of DNA to displace the coordinated Et₄en from its dimethyltin(IV) complex. The nucleotides IMP and GMP have the highest values. The DFT/B3LYP method was used for geometric optimization of the ligand and the complex using the Gaussian 09 program. Also the vibrational frequencies of the ligands and complexes were computed for the optimized geometries. The results shows that there is no imaginary frequencies as found in the calculated vibrational frequencies. The binding energies of the dimethyltin(IV) complexes were calculated. All calculated binding energy values are negative.

Keywords: Dimethyltin(IV); N,N,N',N'-tetraethylethylenediamine; stability constants; DFT calculation.

Resumen. El equilibrio de formación de complejos de dimetilestaño(IV) con N,N,N',N'-tetraetiletilenediamina (Et₄en) fue investigado. La estequiometría y constantes de estabilidad de los complejos formados en solución se determinaron a diferentes temperaturas (15 °C – 35 °C) y en disoluciones de dioxano/agua con diferentes proporciones (15% - 62.5%). El modelo aceptado está constituido de las especies 110, 111, 11-1 y 11-2. Los parámetros termodinámicos ΔH° and ΔS° asociados con la protonación de la N,N,N',N'-tetraetiletilenediamina (Et₄en) y la formación de sus complejos con dimetilestaño(IV) fueron determinados. La reacción de la formación de los complejos es exotérmica. Las constantes de equilibrio del desplazamiento de la N,N,N',N'-tetraetiletilenediamina coordinada al dimetilestaño(IV) con diferentes fragmentos de la ADN fueron calculadas. Los valores K_{eq} indican claramente la capacidad de la ADN de sustituir a la Et₄en de su complejo de dimetilestaño(IV). Los nucleótidos IMP y GMP presentaron los valores más altos. El método DFT/B3LYP fue utilizado para la optimización geométrica del ligante y del complejo utilizando el programa Gaussian 09. Asimismo, se calcularon las frecuencias vibracionales para las geometrías optimizadas de los ligantes y complejos. Los resultados mostraron que todas las frecuencias vibracionales calculadas son reales. También se

calcularon los valores de energías de enlace para los complejos de dimetilestaño(IV) resultando en valores negativos.

Palabras clave: Dimetilestaño(IV); N,N,N',N'-tetraetilethylenediamina; constante de estabilidad; cálculos DFT.

Introduction

The biological activity of organotin(IV) compounds and their use as metallopharmaceuticals has been reported [1-13]. A tremendous research is directed towards the design of non-platinum chemotherapeutics with the aim to optimize the features of classical platinum drugs constituting the basic cisplatin framework viz. their toxic side effects, inherent intrinsic resistance and high cost [14]. Diorganotin(IV) complexes are acting as potential antitumours. They exhibit promise activity against many tumour cell lines [15-19]. Numerous diorganotin(IV) derivatives have been found to exhibit high *in vivo* cytotoxicity against P388 lymphocytic leukaemia.

The antitumour action mechanism is still unclear, although some evidence is consistent with their binding directly to DNA. The antitumour activity of the organotin(IV) complex R_2SnX_2L is depending on the group attached to the tin atom, leaving groups (X) and the coordinated ligand (L). The coordinated ligand favours in some way the transport of the drug into cells, and the antitumour activity would be exerted by the diorganotin(IV) ion released from the complex [20]. The released diorganotin(IV) species would interact with nucleic acids, similarly as in the case of the widely used anticancer drug cisplatin. Therefore, there is a relationship between the stability of the organotin(IV) complexes and their antitumour activity. In conjunction with our previous studies on organotin(IV) complexes [21-26] the present paper aims to study the dimethyltin(IV) complexes with N,N,N',N'-tetraethylethylenediamine to have information regarding the plausible interaction mechanism toward DNA fragment. The four ethyl groups attached to ethylenediamine nitrogen atoms may undergo hydrophobic interaction with the methyl groups of tin(IV) moiety. Such interaction may increase the stability of the formed complex. Also, the four ethyl groups of Et₄en will increase the hydrophobic nature of the complex, which will facilitate the dimethyltin(IV) complex to get into the cell. Density functional theory (DFT) calculations for geometry optimization and vibrational frequencies of all complexes and ligands were reported.

Experimental

Materials and Reagents

Dimethyltin(IV) dichloride (DMT) was obtained from Merck Chem. Co. The ligand N,N,N',N'-tetraethylethylenediamine was supplied by Sigma Chem. Co., 1,4-Dioxane was provided by Aldrich Chem. Co. Sodium hydroxide solutions were prepared by diluting the content of BDH concentrated volumetric solutions vials. NaOH solutions were frequently checked by titration against potassium hydrogen phthalate.

Procedure and Measuring Techniques

A Metrohm 686 titroprocessor was used for the potentiometric titrations. The titroprocessor is equipped with a 665 dosimat (Switzerland-Herisau) and a Metrohm glass-calomel combined electrode is used. NBS standard buffer solutions [27] were used to calibrate the titroprocessor and electrode. The temperature of the titrated solutions was adjusted by circulating thermostatically controlled water through the jacket of a titration vessel. Sodium nitrate was used to adjust the ionic strength to 0.1 mol dm⁻³. All titrations were performed under a nitrogen atmosphere. N,N,N',N'-tetraethylethylenediamine solution was prepared in the protonated form by dissolving in HNO₃ solution. The protonation constants of Et₄en were determined by titrating 1.25 mmol. The hydrolysis constants of dimethyltin(IV) were determined by titrating 1.25 mmol. The formation constants of Et₄en complexes with dimethyltin(IV) were determined by titrating solution mixture of Et₄en (1.25 mmol) and dimethyltin(IV) (1.25 mmol). The volume of each titrated solution mixtures is reached

to 40 mL and the ionic strength was 0.1 mol. dm⁻³ (adjusted with NaNO₃). 0.05 mol. dm⁻³ NaOH solution was used as titrant. The pK_w values of solution mixtures of dioxane-H₂O were estimated as reported previously [28,29]. In this procedure various amounts of standard NaOH solution were added to a 0.10 mol. dm⁻³ NaNO₃ solution. From the amount of base added, [OH⁻] was calculated. [H⁺] was calculated from the pH value. The values obtained in this way for log[OH⁻][H⁺] (log K_w) are -14.23, -14.50, -14.92, -15.12 and -15.63 for 25.0, 37.5, 50.0, 62.5 and 75.0% dioxane in H₂O, respectively.

The titration data were processed to evaluate the equilibrium constants expressed by Eq. 1 and Eq. 2.



$$\beta_{pqr} = \frac{[M_p L_q H_r]}{[M]^p [L]^q [H]^r} \quad 2$$

Where M, L and H stand for dimethyltin(IV), Et₄en and proton respectively. The compound of general formula M_pL_qH_r formed is expressed by the coefficients p, q and r, where M, L and H stand for dimethyltin(IV), Et₄en and proton respectively. The formation constants were evaluated using the computer program MINIQUAD-75 [30]. The stoichiometry and formation constants of the complexes formed were determined by trying various possible models of different composition. The accepted model is giving the best statistical fit and chemically consistent with the magnitude of various residuals, as reported previously [30]. Tables 1 and 2 include the formation constants and their standard deviation as obtained from the MINIQUAD output. The speciation diagrams were evaluated using the program SPECIES [31].

Table 1. Formation constants of dimethyltin (IV) complexes with Et₄en in water at different temperatures.

System ^a	Temp(°C)	p	q	r ^b	Log β ^c	S ^d
DMT	15	1	0	-1	-3.56(0.01)	5.4E-8
		1	0	-2	-9.05(0.01)	
		1	0	-3	-19.79(0.04)	
		1	0	-4	-30.41(0.04)	
		2	0	-2	-4.23(0.01)	
		2	0	-3	-9.52(0.01)	
		2	0	-4	-15.23(0.01)	
DMT-Et ₄ en	15	0	1	1	9.94(0.01)	1.5E-8
		0	1	2	16.80(0.01)	
		1	1	0	11.71(0.03)	
		1	2	0	18.53(0.06)	
		1	1	1	16.69(0.05)	
		1	1	-1	5.32(0.05)	
		1	1	-2	-2.52(0.05)	
DMT	20	1	0	-1	-3.31(0.01)	6.1E-8
		1	0	-2	-8.64(0.01)	
		1	0	-3	-19.27(0.05)	
		1	0	-4	-30.71(0.02)	
		2	0	-2	-3.77(0.01)	
		2	0	-3	-8.91(0.02)	
		2	0	-4	-14.44(0.01)	

DMT-Et ₄ en	20	0	1	1	9.81(0.01)	1.4E-8 3.2E-8
		0	1	2	16.54(0.01)	
		1	1	0	11.52(0.04)	
		1	2	0	18.23(0.05)	
		1	1	1	16.47(0.05)	
		1	1	-1	5.17(0.04)	
		1	1	-2	-2.41(0.04)	
DMT	25	1	0	-1	-3.03(0.01)	4.3E-8
		1	0	-2	-8.21(0.01)	
		1	0	-3	-18.73(0.03)	
		1	0	-4	-29.54(0.02)	
		2	0	-2	-3.12(0.01)	
		2	0	-3	-8.13(0.02)	
		2	0	-4	-13.59(0.02)	
DMT-Et ₄ en	25	0	1	1	9.65(0.01)	3.4E-8 4.1E-10
		0	1	2	16.22(0.02)	
		1	1	0	11.31(0.02)	
		1	2	0	17.80(0.04)	
		1	1	1	16.23(0.04)	
		1	1	-1	5.00(0.04)	
		1	1	-2	-2.24(0.05)	
DMT	30	1	0	-1	-2.81(0.01)	6.6E-8
		1	0	-2	-7.91(0.02)	
		1	0	-3	-18.38(0.05)	
		1	0	-4	-28.71(0.02)	
		2	0	-2	-2.87(0.02)	
		2	0	-3	-7.84(0.04)	
		2	0	-4	-13.06(0.04)	
DMT-Et ₄ en	30	0	1	1	9.51(0.01)	9.3E-9 8.7E-10
		0	1	2	15.94(0.01)	
		1	1	0	11.08(0.03)	
		1	2	0	17.32(0.09)	
		1	1	1	15.98(0.04)	
		1	1	-1	4.81(0.07)	
		1	1	-2	-2.11(0.05)	
DMT	35	1	0	-1	-2.49(0.02)	6.9E-8
		1	0	-2	-7.54(0.02)	
		1	0	-3	-17.95(0.06)	
		1	0	-4	-28.20(0.03)	
		2	0	-2	-2.27(0.03)	
		2	0	-3	-7.11(0.04)	
		2	0	-4	-12.34(0.04)	
DMT-Et ₄ en	35	0	1	1	9.42(0.01)	1.7E-8 1.1E-9
		0	1	2	15.70(0.01)	
		1	1	0	10.90(0.05)	
		1	2	0	16.96(0.10)	
		1	1	1	15.78(0.05)	
		1	1	-1	4.66(0.07)	
		1	1	-2	2.90(0.05)	

^a p, q and r are the stoichiometric coefficient corresponding to organotin(IV), Et₄en and H⁺, respectively; ^c Standard deviations are given in parentheses; ^d Sum of square of residuals.

Table 2. Formation constants of dimethyltin (IV) complexes with Et₄en in dioxane-water solutions of different compositions.

System	Dioxane %	p	q	r ^a	Log β ^b	S ^c
DMT	12.5	1	0	-1	-3.27(0.01)	1.3E-7
		1	0	-2	-8.70(0.02)	
		1	0	-3	-19.53(0.06)	
		1	0	-4	-30.09(0.03)	
		2	0	-2	-3.68(0.02)	
		2	0	-3	-8.94(0.05)	
		2	0	-4	-14.52(0.04)	
DMT-Et ₄ en	12.5	0	1	1	9.75(0.01)	1.8E-8 1.5E-9
		0	1	2	16.40(0.02)	
		1	1	0	11.37(0.02)	
		1	2	0	18.52(0.07)	
		1	1	1	16.00(0.04)	
		1	1	-1	5.00(0.07)	
		1	1	-2	-2.14(0.05)	
DMT	25.0	1	0	-1	-3.39(0.00)	1.0E-8
		1	0	-2	-8.99(0.01)	
		1	0	-3	-19.83(0.01)	
		1	0	-4	-31.30(0.01)	
		2	0	-2	-3.92(0.01)	
		2	0	-3	-9.27(0.04)	
		2	0	-4	-15.11(0.03)	
DMT-Et ₄ en	25.0	0	1	1	9.57(0.00)	2.8E-9 1.5E-9
		0	1	2	16.02(0.01)	
		1	1	0	11.11(0.03)	
		1	2	0	18.10(0.07)	
		1	1	1	15.80(0.04)	
		1	1	-1	4.84(0.10)	
		1	1	-2	-2.31(0.06)	
DMT	37.5	1	0	-1	-3.53(0.01)	1.4E-8
		1	0	-2	-9.23(0.01)	
		1	0	-3	-20.18(0.02)	
		1	0	-4	-31.83(0.01)	
		2	0	-2	-4.17(0.01)	
		2	0	-3	-9.61(0.03)	
		2	0	-4	-15.59(0.03)	
DMT-Et ₄ en	37.5	0	1	1	9.38(0.01)	1.1E-8 1.3E-9
		0	1	2	15.72(0.01)	
		1	1	0	10.72(0.02)	
		1	2	0	17.50(0.02)	
		1	1	1	15.48(0.04)	
		1	1	-1	4.53(0.01)	
		1	1	-2	-2.64(0.04)	
DMT	50.0	1	0	-1	-3.60(0.01)	2.8E-8
		1	0	-2	-9.43(0.01)	
		1	0	-3	-20.58(0.02)	
		1	0	-4	-32.52(0.02)	
		2	0	-2	-4.30(0.01)	
		2	0	-3	-9.78(0.04)	

		2	0	-4	-16.03(0.04)	
DMT-Et ₄ en	50.0	0	1	1	9.17(0.01)	1.5E-8
		0	1	2	15.30(0.01)	
		1	1	0	10.31(0.04)	7.5E-10
		1	2	0	16.90(0.07)	
		1	1	1	15.13(0.06)	
		1	1	-1	4.25(0.09)	
		1	1	-2	-2.94(0.06)	
DMT	62.5	1	0	-1	-3.68(0.01)	8.7E-8
		1	0	-2	-9.62(0.02)	
		1	0	-3	-20.89(0.04)	
		1	0	-4	-33.26(0.05)	
		2	0	-2	-4.47(0.01)	
		2	0	-3	-9.90(0.03)	
		2	0	-4	-16.51(0.04)	
DMT-Et ₄ en	62.5	0	1	1	9.04(0.01)	1.1E-8
		0	1	2	15.11(0.01)	
		1	1	0	9.93(0.03)	7.4E-10
		1	2	0	16.29(0.09)	
		1	1	1	14.82(0.05)	
		1	1	-1	3.96(0.07)	
		1	1	-2	-3.26(0.06)	

^ap, q and r are the stoichiometric coefficient corresponding to organotin(IV), Et₄en and H⁺ respectively; ^bStandard deviations are given in parentheses; ^cSum of square of residuals.

Computational Details

For geometries optimization, the density functional theory was applied using the Gaussian09 program [32]. A full optimization of all complexes and ligands geometries were obtained using the hybrid functional B3LYP (Becke's, three-parameter exchange functional, in combination with Lee-Yang-Parr correlation function) [33-35]. The basis set used is LANL2DZ (Los Alamos National Laboratory 2 double-zeta) [36-38] for Sn metal, while C, H, N, O and Cl were described by 6-311G* standard basis set in optimization. The GaussView 5.0 by Gaussian Inc. program was used for inspecting the input and output files generated by Gaussian09, for preprocessing, structure modification, and post processing analyses of structures, frequencies, and forces. To identify the most stable structure, the minima, a frequency analysis was performed for each stationary point. These analyses are performed to ensure that all minima have no imaginary frequencies in the vibrational mode calculations.

Results and discussion

Dimethyltin(IV) complex formation equilibria

The protonation constants of N,N,N',N'-tetraethylmethylenediamine were determined using the same experimental conditions as solvent composition, temperature and ionic strength which are used for the investigation of organotin complex equilibria. The overall protonation constants ($\log\beta_{11}$ and $\log\beta_{12}$) for Et₄en were estimated. The values obtained agree with the literature data [39] considering changes in experimental condition.

The acid-base equilibria of dimethyltin(IV) ion was investigated previously in aqueous media [40-44]. The potentiometric data in the mixed solvents used were fitted assuming a model composed of the species 10-1, 10-2, 10-3, 10-4, 10-4, 20-3 and 20-4, Table 1. The aqua-hydroxo species (10-1) may undergo dimerization according to the general equilibrium




The dimerization constant (K_d) can be determined by Eq. 4.

$$\log K_d = \log \beta_{202-} - 2\log \beta_{10-1} \quad 4$$

Following the same way the dimerization constant of the species 20-4 is determined by the Eq. 5 [24].

$$\log K_d = \log \beta_{20-4} - 2\log \beta_{10-2} \quad 5$$

The potentiometric titration curves of the protonated N,N,N',N'-tetraethylethylenediamine and its dimethyltin(IV) complex are compared. The complex titration curve is lower than the Et₄en curve, Fig. 1. This corresponds to the formation of a complex species through the ionization of hydrogen ion. The titration data were fitted considering the formation of the 110, 111, 11-1 and 11-2 species, according to scheme 1.



Scheme 1. Formation equilibria of dimethyltin(IV)-Et₄en complex .

Eq. 6 and 7 [26] are used to calculate the pK_a of coordinated water molecules.

$$pK_{a1} = \log \beta_{110-} - \log \beta_{11-1} \quad 6$$

$$pK_{a2} = \log \beta_{11-1} - \log \beta_{11-2} \quad 7$$

The obtained values are 5.57 and 9.68 for pK_{a1} and pK_{a2} respectively. These are higher than those of water molecules coordinated to the free dimethyltin(IV) ion. This may be explained on the premise that

coordination of Et₄en will decrease the electrophilicity of tin. This will result in the coordinated water molecule will be weakly bound and consequently will be less acidic.

The speciation diagram of DMT-Et₄en complexes is used to provide a useful picture of the coordination of organotin with Et₄en as a function of pH, Fig. 1. The protonated complex with stoichiometric coefficients 111, predominates at lower pH. It attains a maximum concentration of 38% at pH = 3.8. The deprotonated complex 110 starts to form at pH 3.0 and reaches the maximum concentration of 39% at pH = 5.7. The hydrolyzed form 11-1 predominates between pH = 5.7 and 8.0 with maximum formation degree of 27% at pH = 6.9. It is interesting to note that the complex species 110 and 11-1 are predominating in the physiological pH range. Therefore the binding of DMT complex with DNA constituent, the main target in the chemotherapy is possible. This may be explained on the premise that the DNA constituents coordinate in the negatively charged anionic form.




Fig. 1. Concentration distribution of various species as a function of pH in the DMT- Et₄en system.

Effect of temperature

The thermodynamic parameters ΔH° and ΔS° associated with the protonation of N,N,N',N'-tetraethylenthiediamine (Et₄en) and its complex formation with the dimethyltin(IV) species were calculated from the temperature dependence of the data in Table 3. The plots are given in figures 2-3. These data can be employed to extrapolate the equilibrium constants to other temperatures. The main conclusions from the data can be summarized as follows:

a) The protonation reactions (8) and (10) of Et₄en are exothermic and of comparable ΔH° value. Three factors affect the protonation reactions (i). The neutralization reaction, which is an exothermic process. (ii). Desolvation of ions, which is an endothermic process. (iii). The change of the conformation and the arrangements of the hydrogen bonds around the free and the protonated ligands.

b) The complexation reaction (11-15) for dimethyltin(IV) ion with Et₄en is exothermic with negative ΔH° values, amounting to -70.0, -137.5, -78.5 and -57.1 kJ.mol⁻¹.

Table 3. Thermodynamic parameters for the equilibria of dimethyltin(IV) complexes.^a

Equilibrium	ΔH° (kJmol ⁻¹)	ΔS° (JK ⁻¹ mol ⁻¹)
DMT ^b		
1) $M(H_2O)_4^{2+} + OH^- \rightleftharpoons M(H_2O)_3(OH)^+ + H_2O$	40.2(0.5)	344(2)
2) $M(H_2O)_3(OH)^+ + OH^- \rightleftharpoons M(H_2O)_2(OH)_2 + H_2O$	-11.5(0.2)	128(1)
3) $M(H_2O)_2(OH)_2 + OH^- \rightleftharpoons M(H_2O)(OH)_3^- + H_2O$	-21.4(0.4)	-7(1)

4) $M(H_2O)(OH)_3^- + OH^- \rightleftharpoons M(OH)_4^{2-} + H_2O$	-20.9(0.4)	-4(1)
5) $2M(H_2O)_4^{2+} + 2OH^- \rightleftharpoons M_2(H_2O)_4(OH)_2^{2+} + 4H_2O$	65.0(0.8)	690(7)
6) $M_2(H_2O)_4(OH)_2^{2+} + OH^- \rightleftharpoons M_2(H_2O)_3(OH)_3^+ + H_2O$	-13.0(0.3)	127(1)
7) $M_2(H_2O)_3(OH)_3^+ + OH^- \rightleftharpoons M_2(H_2O)_2(OH)_4^+ + H_2O$	-6.1(0.1)	142(1)
Et ₄ en		
8) $L + H^+ \rightleftharpoons LH^+$	-45.6(0.1)	32.0(0.3)
9) $LH^+ + H^+ \rightleftharpoons LH_2^{2+}$	-45.78	31.98
10) $L + 2H^+ \rightleftharpoons LH_2^{2+}$	-95.2(0.1)	-8.6(0.4)
DMT-Et ₄ en		
11) $M + L \rightleftharpoons ML$	-70.0(0.1)	-18.6(0.3)
12) $M + 2L \rightleftharpoons ML_2$	-137.5(0.3)	-121.4(1)
13) $M + L + H \rightleftharpoons MLH$	-78.5(0.1)	47.1(0.3)
14) $M + L - H \rightleftharpoons MLH_{-1}$	-57.1(0.1)	-96.0(0.3)
15) $M + L - 2H \rightleftharpoons MLH_{-2}$	39.5(0.2)	89.2(0.8)

^aM denotes $(CH_3)_3Sn^{2+}$; L denote; Standard deviations are given in parentheses. ^bhydrolysis data are from reference [51]




Fig. 2. Plot of $\log \beta_1$ vs. $1/T$ for $(CH_3)_2Sn$ -Et₄en complex.




Fig. 3. Plot of $\log \beta_2$ vs. $1/T$ for $(\text{CH}_3)_2\text{Sn}\text{-Et}_4\text{en}$ complex.

Effect of solvent

It is reported that the effective dielectric constant of the surrounding of active sites cavities of enzymes and proteins [45,46] is smaller, compared to in bulk water. Dielectric constants in such sites were found to be in range from 30 to 70 [47, 48]. Hence by using solution media containing around 10 to 50% dioxane, one may expect to approach to some extent the situation in these biological sites. By this way one may extrapolate the data to physiological condition. Careful examination of the medium effect on the equilibrium constants reveal the following features:

- pK_a of Et₄en decrease linearly with increasing the percentage of organic solvent in the medium. This behavior is discussed based on the hydrophobic interaction between the neutral Et₄en species and dioxane.
- The pK_a values of coordinated water molecules in $[(\text{CH}_3)_2\text{Sn}(\text{OH}_2)_4]^{2+}$ increase with increasing the dioxane percentage in the medium. This may be explained on the premise that the solvent of relatively low dielectric constant will increase the electrostatic forces between proton and hydrolyzed form of DMT.
- The formation constants for dimethyltin(IV) complexes with Et₄en decrease upon addition of dioxane to an aqueous solution of the corresponding species, figures 4-5. This can be explained by better solvation of virtually hydrophobic species $\text{CH}_3\text{Sn}^+/\text{CH}_3\text{SnCl}$ by dioxane through formation of coordinative bonds between O-dioxane and Sn centres. This leads to stabilization and modulation of solubility of dimethyltin(IV) species. This results in lowering the complex stability. This finding is in agreement with that proposed for alkyltin(IV) complexes with D-glucosamine [49].




Fig. 4. Plot of $\log \beta_1$ vs. dioxane percentage for $(\text{CH}_3)_2\text{Sn-Et}_4\text{en}$ complex.





Fig. 5. Plot of $\log \beta_2$ vs. dioxane percentage for $(\text{CH}_3)_2\text{Sn-Et}_4\text{en}$ complex.

Displacement reaction of coordinated Et₄en by DNA constituents


It was reported that $(CH_3)_2Sn(IV)$ has high affinity for N-donor ligands such as DNA constituents. Such interaction with DNA is thought to be responsible for the anti-tumour activity of related complexes [50]. The mechanism of antitumor activity of $(CH_3)_2Sn$ -amine complex is based on the displacement of the amine by DNA. Consequently, the equilibrium constant for such reaction is of biological importance. Consider inosine as a typical DNA constituent (presented by HA) and Et₄en (presented by B). The equilibria of complex-formation and displacement reactions may be presented introductory as:



$$\beta_{110} (\text{CH}_3)_2\text{Sn}(\text{A})^+ = [(\text{CH}_3)_2\text{Sn}(\text{A})^+] / [(\text{CH}_3)_2\text{Sn}^{2+}] [\text{A}^-] \quad 8b$$



$$\beta_{110} (\text{CH}_3)_2\text{SnB}^{2+} = [(\text{CH}_3)_2\text{SnB}^{2+}] / [(\text{CH}_3)_2\text{Sn}^{2+}][\text{B}] \quad 9\text{b}$$



The equilibrium constant for the displacement reaction is expressed by:

$$K_{eq} = [(CH_3)_2Sn(A^+)[B]/[(CH_3)_2Sn(B)^{2+}[A^-] \quad 10c$$

Substitution results in:

11

$$K_{eq} = \beta_{110} [(CH_3)_2Sn(A)]^+ / \beta_{110} [(CH_3)_2Sn(B)]^{2+}$$

12

The formation constant value for Et₄en complex, $[(\text{CH}_3)_2\text{SnB}]^{2+}$, amounts to $\log \beta_{110} = 11.31$ at 25°C (Table 1). The $\log \beta_{110}$ values [51] for (CH_3)₂Sn-DNA constituents complexes $[(\text{CH}_3)_2\text{SnA}]^+$ together with the equilibrium constants values ($\log K_{\text{eq}}$) of displacement reaction calculated by Eq. 12, are given in Table 4. The K_{eq} values clearly indicate the ability of DNA to displace the coordinated Et₄en from its dimethyltin(IV) complex. Also, K_{eq} value reveals to what extent the organotin-amine complex could interact with the DNA constituents, the main target in chemotherapy of tumours. The values of equilibrium constant of the displacement reaction for the different DNA constituents are compared. The nucleotide IMP and GMP have the highest values. The extra stability is due to the different columbic force of attraction between the dipositively charged dimethyltin(IV) ion and the nucleotides IMP or GMP having extra negative charges on the phosphate group. Also, the phosphate group of IMP and GMP is acting as anchoring site for tin [52]. This will facilitate the release of coordinated Et₄en.

Table 4. Equilibrium constants for displacement reaction of coordinated N,N,N',N'-tetramethylethylenediamine by DNA constituents at 25°C and 0.1M NaNO₃.

DNA	$\log \beta_{110}^{\text{a}}$ DMT-DNA	$\log K_{\text{eq}}$
Inosine	8.13	-3.18
Inosine-5'-monophosphate	11.90	0.59
Guanosine-5'- monophosphate	12.34	1.03
Adenine	10.01	-1.30
Adenosine	4.41	-6.90
Adenosine -5'-monophosphate	6.07	-5.24
Uracil	9.34	-1.97
Thymine	9.61	-1.70
Thymidine	9.25	-2.06
Cytosine	4.44	-6.87
Cytidine	3.77	-7.54

^ataken from ref. [51]

Geometry optimiation

Calculations of the equilibrium geometries of the complexes and the ligands were carried out by the program Gaussian 09 at the B3LYP level of theory using the bases set LANL2DZ for Sn metal and standard basis set 6-311G* for C, H, N, O and Cl atoms. The optimized structures of the ligands and the complexes are given in Fig. 6. The geometric computational parameters for the optimized molecular geometry as bond angles, atomic charges and bond lengths are determined and given in (Tables 5 and 6). The results are interpreted as given in the following:

- 1- The N-CH₂ bond length in the complexes $((\text{CH}_3)_2\text{Sn}(\text{Et}_4\text{en})\text{Cl}_2$, $(\text{CH}_3)_2\text{Sn}(\text{en})(\text{H}_2\text{O})_2^{2+}$, $(\text{CH}_3)_2\text{Sn}(\text{Et}_4\text{en})(\text{H}_2\text{O})_2^{2+}$ and $(\text{CH}_3)_2\text{Sn}(\text{Et}_4\text{en})(\text{OH})_2$) are 1.50, 1.52 and 1.50 Å respectively. It is longer than the N-CH₂ bond in the free ligand (1.47 Å). This elongation is due to the bonding of the Nitrogen atom with the metal.
- 2- The Sn-N bond length in the $(\text{CH}_3)_2\text{Sn}(\text{Et}_4\text{en})\text{Cl}_2$ complex (2.53-2.54 Å) is longer than that in $(\text{CH}_3)_2\text{Sn}(\text{en})\text{Cl}_2$ complex (2.40 Å). The main reason of this attributed to the changing the charge density on the nitrogen atom (-0.59 on N in en compared to -0.10 on N in Et₄en).
- 3- The bond angle Et₂N-C-C in Et₄en is decreased from (115.8 °) upon complexation to 113.4 ° ($((\text{CH}_3)_2\text{Sn}(\text{Et}_4\text{en})\text{Cl}_2$), 111.8 ° ($((\text{CH}_3)_2\text{Sn}(\text{Et}_4\text{en})(\text{H}_2\text{O})_2^{2+}$)), 113.6 ° ($((\text{CH}_3)_2\text{Sn}(\text{Et}_4\text{en})(\text{OH})_2$))). This may be due to the formation of cyclic compounds.
- 4- The H₂O-Sn(IV) bond length is 2.43-2.47 Å in $(\text{CH}_3)_2\text{Sn}(\text{Et}_4\text{en})(\text{H}_2\text{O})_2^{2+}$ species while 2.39 Å in $(\text{CH}_3)_2\text{Sn}(\text{en})(\text{H}_2\text{O})_2^{2+}$ species. After deprotonation of coordinated water, the O-Sn(IV) bond is

decreased to 2.00 Å for the two species. This is attributed to the strong bond between the negatively charged OH⁻ ion and Sn(IV).

Table 5. Calculated N–C, Sn–N and Sn–O bond lengths of the optimized structures of the ligands and complexes according to B3LYP/LanL2DZ level.

Spices	<i>d</i> _{(N–C), Å}		<i>d</i> _{(Sn–N), Å}		<i>d</i> _{(Sn–O), Å}	
Et ₄ en	N(2)-C(6)	1.47	-		-	
	N(1)-C(3)	1.47				
En	N(2)-C(6)	1.47	-		-	
	N(1)-C(3)	1.47				
(CH ₃) ₂ Sn(Et ₄ en)Cl ₂	N(12)-C(14)	1.50	Sn(1)-N(13)	2.59	-	
	N(13)-C(17)	1.50	Sn(1)-N(12)	2.63		
(CH ₃) ₂ Sn(en)Cl ₂	N(13)-C(17)	1.49	Sn(1)-N(13)	2.40	-	
	N(12)-C(14)	1.49	Sn(1)-N(12)	2.40		
(CH ₃) ₂ Sn(Et ₄ en)(H ₂ O) ₂ ²⁺	N(11)-C(15)	1.52	Sn(1)-N(11)	2.33	Sn(1)-O(48)	2.46
	N(10)-C(12)	1.52	Sn(1)-N(10)	2.33	Sn(1)-O(46)	2.43
(CH ₃) ₂ Sn(en)(H ₂ O) ₂ ²⁺	N(11)-C(15)	1.52	Sn(1)-N(10)	2.28	Sn(1)-O(22)	2.37
	N(10)-C(12)	1.52	Sn(1)-N(11)	2.28	Sn(1)-O(24)	2.37
(CH ₃) ₂ Sn(Et ₄ en)(OH) ₂	N(11)-C(15)	1.50	Sn(1)-N(10)	2.76	Sn(1)-O(48)	1.99
	N(10)-C(12)	1.50	Sn(1)-N(11)	2.83	Sn(1)-O(46)	1.99
(CH ₃) ₂ Sn(en)(OH) ₂	N(10)-C(12)	1.49	Sn(1)-N(10)	2.48	Sn(1)-O(24)	1.99
	N(11)-C(15)	1.49	Sn(1)-N(11)	2.48	Sn(1)-O(22)	1.99

Table 6. Calculated N–C–C bond angles and the nitrogen atom Mulliken charges of the optimized structures of the ligands and complexes according to B3LYP/LanL2DZ level.

Spices	N–C–C bond angle		Nitrogen atom charge	
Et ₄ en	C(6)-C(3)-N(1)	115.5	N(1)	-0.10
	C(3)-C(6)-N(2)	115.9	N(2)	-0.10
En	C(6)-C(3)-N(1)	110.9	N(1)	-0.63
	C(3)-C(6)-N(2)	110.9	N(2)	-0.63
(CH ₃) ₂ Sn(Et ₄ en)Cl ₂	C(14)-C(17)-N(13)	113.9	N(12)	-0.26
	C(17)-C(14)-N(12)	113.1	N(13)	-0.26
(CH ₃) ₂ Sn(Et ₄ en)(H ₂ O) ₂ ²⁺	C(12)-C(15)-N(11)	111.9	N(10)	-0.39
	C(15)-C(12)-N(10)	110.9	N(11)	-0.39
(CH ₃) ₂ Sn(en)Cl ₂	C(17)-C(14)-N(12)	109.5	N(12)	-0.69
	C(14)-C(17)-N(13)	109.5	N(13)	-0.69
(CH ₃) ₂ Sn(en)(H ₂ O) ₂ ²⁺	C(12)-C(15)-N(11)	108.8	N(10)	-0.76
	C(15)-C(12)-N(10)	108.8	N(11)	-0.76
(CH ₃) ₂ Sn(Et ₄ en)(OH) ₂	C(12)-C(15)-N(11)	114.7	N(10)	-0.22
	C(15)-C(12)-N(10)	113.5	N(11)	-0.20
(CH ₃) ₂ Sn(en)(OH) ₂	C(15)-C(12)-N(10)	110.2	N(10)	-0.66
	C(12)-C(15)-N(11)	110.2	N(11)	-0.66




Fig. 6. The optimized structure of ligands and complexes according to B3LYP level of theory using the bases set LANL2DZ for Sn metal and standard basis set 6-311G* for C, H, N, O and Cl atoms.

Based on the DFT calculation, the polarity of the complexes, is much larger than that for the free ligands as indicated by the dipole moment of the complexes and the ligands (Table 6). On the other hand, the dipole moment of the hydroxo-complexes $(\text{CH}_3)_2\text{Sn}(\text{Et}_4\text{en})(\text{OH})_2$ and $(\text{CH}_3)_2\text{Sn}(\text{en})(\text{OH})_2$ are lower than that of the corresponding dichloro-complexes $(\text{CH}_3)_2\text{Sn}(\text{Et}_4\text{en})\text{Cl}_2$ (Table 7). This may be described on the assumption that the O-Sn(IV) bond is shorter than the Cl-Sn(IV) bond as given in Table 5.

The binding energy (BE) of each complex was calculated according to the following equation:

$$\text{BE} = E_{\text{AB}} - (E_{\text{A}} + E_{\text{B}})$$

13

In equation 13, E_{AB} , E_{A} and E_{B} is the energy of the complex, dimethyltin(IV) species and ligand respectively. The calculated BE (in kJ/mol) of the complexes $(\text{CH}_3)_2\text{Sn}(\text{Et}_4\text{en})\text{Cl}_2$, $(\text{CH}_3)_2\text{Sn}(\text{en})\text{Cl}_2$, $(\text{CH}_3)_2\text{Sn}(\text{Et}_4\text{en})(\text{OH})_2$, $(\text{CH}_3)_2\text{Sn}(\text{en})(\text{OH})_2$, $(\text{CH}_3)_2\text{Sn}(\text{Et}_4\text{en})(\text{H}_2\text{O})_2^{2+}$ and $(\text{CH}_3)_2\text{Sn}(\text{en})(\text{H}_2\text{O})_2^{2+}$ are -52.4, -99.2, -4.2, -31.0, -400.1 and -382.8, respectively. The negative values of the BE indicate that the complexes are stable.

Table 7. The calculated energies and dipole moments for of the ligands and complexes according to B3LYP/LanL2DZ level.

Spices	E ^a	HOMO	LUMO	ΔE	Dipole moment
Et ₄ en	-504.9	-4.84	2.76	7.60	0.69
En	-190.5	-5.51	2.70	8.21	1.05
$(\text{CH}_3)_2\text{Sn}(\text{Et}_4\text{en})\text{Cl}_2$	-618.1	-6.54	-0.11	6.43	12.38
$(\text{CH}_3)_2\text{Sn}(\text{en})\text{Cl}_2$	-303.7	-6.64	0.08	6.72	12.47
$(\text{CH}_3)_2\text{Sn}(\text{Et}_4\text{en})(\text{OH})_2$	-739.8	-5.90	0.90	6.80	6.54
$(\text{CH}_3)_2\text{Sn}(\text{en})(\text{OH})_2$	-425.4	-5.88	1.39	7.27	7.33

^a E: is the total energy (a.u.), HOMO: is the highest occupied molecular orbital (eV), LUMO: is the lowest unoccupied molecular orbital (eV), ΔE: E_{LUMO} - E_{HOMO} (eV) and dipole moment calculated (Debye).

Calculated infrared spectra of ligands and complexes

The experimental counterparts of the calculated IR spectra results are currently not available for comparison and it will be considered in our future work. Therefore, we will discuss the calculated IR spectra of the complexes from the theoretical point of view only. The calculated IR spectra of the free ligands and the complexes are shown in Fig. 7. There are no imaginary vibrational frequencies observed in all calculated spectra of the ligands and complexes. The assignments of the calculated vibrational frequencies of the free ligands and their complexes are listed in Table 8. The theoretical assignments are easily obtained by the visualization of the normal mode displacement vectors utilizing the GaussView program. The assignments listed in Table 8 are only for the vibrational modes of the amino group vibrations because they are the most sensitive to changes upon the metal-nitrogen interactions. Also, the Sn-N, Sn-OH, Sn-Cl and Sn-OH₂ stretching vibrations are identified. In the spectra of the $(\text{CH}_3)_2\text{Sn}(\text{en})\text{Cl}_2$, $(\text{CH}_3)_2\text{Sn}(\text{en})(\text{H}_2\text{O})_2^{2+}$ and $(\text{CH}_3)_2\text{Sn}(\text{en})(\text{OH})_2$ complexes the $\beta_s(\text{NH}_2)$ vibration is shifted to higher wave frequency in comparison with that found in the free en ligand (Table 8). Also for the same complexes the $\nu(\text{C}-\text{N})$ vibration mode is shifted to lower wave frequency in comparison with that of free en ligand. However, in the case of $(\text{CH}_3)_2\text{Sn}(\text{Et}_4\text{en})\text{Cl}_2$, $(\text{CH}_3)_2\text{Sn}(\text{Et}_4\text{en})(\text{H}_2\text{O})_2^{2+}$ and $(\text{CH}_3)_2\text{Sn}(\text{Et}_4\text{en})(\text{OH})_2$ complexes this vibrational mode ($\nu(\text{C}-\text{N})$) is shifted to higher frequency in comparison with that of free Et₄en ligand. In the spectra of all complexes the $\nu_{as}(\text{C}-\text{C}-\text{N}$ vibration mode is shifted to lower frequency in comparison with those of free en and Et₄en ligands. The above mentioned shifts of these vibrational modes can be attributed to the coordination of the nitrogen atom with the metal.

Table 8. Calculated infrared absorption (cm^{-1}) data for the ligands and complexes.

Compounds	$\nu(\text{Sn}-\text{Cl})$	$\nu(\text{Sn}-\text{OH})$	$\nu(\text{Sn}-\text{OH}_2)$	$\nu(\text{Sn}-\text{N})$	$\nu_{as}(\text{Sn}-\text{CH}_3)$	$\nu_s(\text{Sn}-\text{CH}_3)$	$\beta_s(\text{NH}_2)$	$\nu(\text{C}-\text{N})$	$\nu_{as}(\text{C}-\text{C}-\text{N})$
Et ₄ en	-	-	-	-	-	-	-	1021	1157
En	-	-	-	-	-	-	1670	1066	1172
$(\text{CH}_3)_2\text{Sn}(\text{Et}_4\text{en})\text{Cl}_2$	261	-	-	576 612	537	471	-	1035	1133

$(CH_3)_2Sn(en)Cl_2$	260		-	285	552	490	1696	1027	1135
$(CH_3)_2Sn(Et_4en)(H_2O)_2^{2+}$	-	-	246 242	510 594	579	499	-	1036	1136
$(CH_3)_2Sn(en)(H_2O)_2^{2+}$	-	-	244 249	331	555	512	1696	1012	1064
$(CH_3)_2Sn(Et_4en)(OH)_2$	-	519 524	-	568 603	541	466	-	1067	1131
$(CH_3)_2Sn(en)(O)H_2$	-	524 527	-	355	534	475	1691	1006	1131

^a: Calculated value according to B3LYP/LANL2DZ level, n_s: symmetric stretching, n_{as}: asymmetric stretching n: stretching, β_s: scissoring




Fig. 7. Calculated Infrared spectra of the (a) Et₄en ligand, (b) En ligand, (c) $(CH_3)_2Sn(Et_4en)Cl_2$ complex, (d) $(CH_3)_2Sn(en)Cl_2$ complex, (e) $(CH_3)_2Sn(Et_4en)(H_2O)_2^{2+}$ complex, (f) $(CH_3)_2Sn(en)(H_2O)_2^{2+}$ complex, (g) $(CH_3)_2Sn(Et_4en)(OH)_2$ complex and (h) $(CH_3)_2Sn(en)(OH)_2$ complex according to B3LYP level of theory using the bases set LANL2DZ for Sn metal and standard basis set 6-311G* for C, H, N, O and Cl atoms.

Conclusion

The present investigation reports the complex formation equilibria of dimethyltin(IV) with N,N,N',N'-tetraethylethylenediamine and its parent ethylenediamine. The results show the formation of 1:1 complex and the mono- and dihydroxo- species. The effect of solvent polarity on the stability of the complexes was discussed. The thermodynamic parameters of the complexes as ΔH° and ΔS° were determined and discussed. The displacement reaction of Et₄en coordinated to dimethyltin (IV) by DNA constituents was investigated. IMP and GMP have the highest tendency to displace Et₄en. The geometric parameters, bond angle, atomic charge and bond lengths were computed and interpreted. The results of the present study are expected to contribute to the chemistry of diorganotin (IV) as potential antitumours.

References

1. Gasser, G.; Ott, I.; Metzler-Nolte, N. *J. Med. Chem.* **2011**, *54*, 3-25 DOI: <https://doi.org/10.1021/jm100020w>
2. Gianferrara, T.; Bratsos, I.; Alessio, E. *Dalton Trans.* **2009**, 7588-7598 DOI: <https://doi.org/10.1039/b905798f>
3. Moody, L.; Holder, A. A. *Annu. Rep. Prog. Chem., Sect. A* **2009**, *105*, 505-524 DOI: <https://doi.org/10.1039/b818288b>
4. Hadjikakou, S.K.; Hadjiliadis, N. *Coord. Chem. Rev.* **2009**, *253*, 235-249 DOI: <https://doi.org/10.1016/j.ccr.2007.12.026>
5. Tabassum, S.; Pettinari, C. *J. of Organomet. Chem.* **2006**, *691*, 1761-1766 DOI: <https://doi.org/10.1016/j.jorgchem.2005.12.033>
6. Pellerito, C.; Nagy, L.; Pellerito, L.; Szorescik, A. *J. of Organomet. Chem.* **2006**, *691*, 1733-174 DOI: <https://doi.org/10.1016/j.jorgchem.2005.12.025>
7. Gielen, M.; Biesemans, M.; Willem, R. *Appl. Organometal. Chem.* **2005**; *19*, 440-450 DOI: <https://doi.org/10.1002/aoc.771>
8. Pellerito, L.; Nagy, L. *Coord. Chem. Rev.* **2002**, *224*, 111-150 DOI: [https://doi.org/10.1016/S0010-8545\(01\)00399-X](https://doi.org/10.1016/S0010-8545(01)00399-X)
9. Clarke, M. J.; Zhu, F.; Frasca, D. R. *Chem. Rev.* **1999**, *99*, 2511-2533 DOI: <https://doi.org/10.1021/cr9804238>
10. Gielen, M. *Coord. Chem. Rev.* **1996**, *151*, 41-51.
11. Sadler, P.J. *Adv. Inorg. Chem.* **1991**, *36*, 1-48 DOI: [https://doi.org/10.1016/S0898-8838\(08\)60035-5](https://doi.org/10.1016/S0898-8838(08)60035-5)
12. Guo, Z.; Sadler, P.J. *Angew. Chem. Int. Ed.* **1999**, *38*, 1512-1531 DOI: [https://doi.org/10.1002/\(SICI\)1521-3773\(19990601\)38:11<1512::AID-ANIE1512>3.0.CO;2-Y](https://doi.org/10.1002/(SICI)1521-3773(19990601)38:11<1512::AID-ANIE1512>3.0.CO;2-Y)
13. Tlahuext-Aca, A.; Höpfel, H.; Medrano-Valenzuela, F.; Guerrero-Alvarez, J.; Tlahuext, H.; Ochoa Lara, K.; Reyes-Márquez, V.; Tlahuextl, M. *Z. Anorg. Allg. Chem.* **2012**, *638*, 1-9
14. Kotsakis, A.; Hatzidakis, D.; Vamvakas, L.; Vardakis, N.; Kalykaki, A.; Bozionelou, V.; Androulakis, N.; Kalbakis K.; Saridaki, Z.; Georgoulias, V. *Anticancer Res.*, **2010**, *30*, 4335-4342 DOI: <https://doi.org/10.1002/zaac.201200099>
15. Gielen, M.; Willem, R. *Anticancer Research*, **1992**, *12*, 269-271.
16. Crowe, A. J. *Antitumour activity of tin compounds*, in: *Metal Compounds in Cancer Therapy*, Springer, **1994**, 147-179.
17. Crowe, A. J.; Smith, P. J.; Cardin, C. J.; Parge, H. E.; Smith, F. E. *Cancer Lett.* **1984**, *24*, 45-48 DOI: [https://doi.org/10.1016/0304-3835\(84\)90078-8](https://doi.org/10.1016/0304-3835(84)90078-8)

18. Gielen, M.; Meriem, A.; Boualam, M.; Willem, R. *In vivo (Athens, Greece)* **1993**, *7*, 171-174.
19. Yahyi, A.; Et-Touhami, A.; Yahyaoui, R.; Touzani, R. *Lett. Drug Des. Discovery* **2010**, *7*, 534-540.
20. Muhammad, N.; Shah, A.; Shuja, S.; Ali, S.; Qureshi, R.; Meetsma, A.; Tahir, M.N. *J. Organomet. Chem.* **2009**, *694*, 3431-3437 DOI: <https://doi.org/10.1016/j.jorganchem.2009.06.036>
21. Mohamed, M. M. A.; Shehata, M. R.; Shoukry, M. M. *J. Coord. Chem.* **2001**, *53*, 125-142 DOI: <https://doi.org/10.1080/00958970108022607>
22. Al Alousi, A. S.; Shehata, M. R.; Shoukry, M. M.; Mohamed, N. M. *Chem. Speciation Bioavailability* **2009**, *21*, 1-6 DOI: <https://doi.org/10.3184/095422909X416216>
23. Al-Najjar, A.; Mohamed, M. M. A.; Shehata, M. R.; Shoukry, M. M. *Ann. Chim. (Rome, Italy)* **2006**, *96*, 97-107 DOI: <https://doi.org/10.1002/adic.200690010>
24. Al-Najjar, A.; Mohamed, M. M. A.; Shoukry, M. M. *J. of Coord. Chem.* **2006**, *59*, 193-206 DOI: <http://dx.doi.org/10.1080/00958970500270786>
25. Mohamed, M. M. A.; Shoukry, M. M. *Chem. Pharm. Bull.* **2001**, *49*, 253-257 DOI: <https://doi.org/10.1248/cpb.49.253>
26. Shehata, M. R.; Mohamed, M. M. A.; Shoukry, M. M.; Hussein, M. A.; Hussein, F. M. *J. Coord. Chem.* **2015**, *68*, 1101-1114 DOI: <http://dx.doi.org/10.1080/00958972.2015.1007962>
27. Bates, R. *Theory and Practice* Wley, New York, Sydney (1964).
28. Motekaitis, R. J.; Martell, A. E.; Nelson, D. A. *Inorg. Chem.* **1984**, *23*, 275-283 DOI: <https://doi.org/10.1021/ic00171a005>
29. Shoukry, M. M.; Shehata, M. R.; Hamza, M. S.; van Eldik, R. *Dalton Trans.* **2005**, 3921-3926 DOI: <http://dx.doi.org/10.1039/b510553f>
30. Gans, P.; Sabatini, A.; Vacca, A. *Inorg. Chim. Acta* **1976**, *18*, 237-239 DOI: [https://doi.org/10.1016/S0020-1693\(00\)95610-X](https://doi.org/10.1016/S0020-1693(00)95610-X)
31. L. Pettit, *Personal communication*, 1993.
32. Robb, M. A.; Cheeseman, J. R.; Scalmani, G.; Barone, V.; Mennucci, B.; Petersson, G. A.; Nakatsuji, H.; Caricato, M.; Li, X.; Hratchian, H. P.; Izmaylov, A. F.; Bloino, G. Z. J.; Sonnenberg, J. L.; Hada, M.; Ehara, M.; Toyota, K.; Fukuda, R.; Hasegawa, J.; Ishida, M.; Nakajima, T.; Honda, Y.; Kitao, O.; Nakai, H.; Vreven, T.; Montgomery, J. A.; Peralta, Jr. J. E.; Ogliaro, F.; Bearpark, M.; Heyd, J. J.; Brothers, E.; Kudin, K. N.; Staroverov, V. N.; Kobayashi, R.; Normand, J.; Raghavachari, K.; Rendell, A.; Burant, J. C.; Iyengar, S. S.; Tomasi, J.; Cossi, M.; Rega, N.; Millam, J. M.; Klene, M.; Knox, J. E.; Cross, J. B.; Bakken, V.; Adamo, C.; Jaramillo, J.; Gomperts, R.; Stratmann, R. E.; Yazyev, O.; Austin, A. J.; Cammi, R.; Pomelli, C.; Ochterski, J. W.; Martin, R. L.; Morokuma, K.; Zakrzewski, V. G.; Voth, G. A.; Salvador, P.; Dannenberg, J. J.; Prich, S. D.; Daniels, A. D.; Farkas, O.; Foresman, J. B.; Ortiz, J. V.; Cioslowski, J.; Fox, D. J. Wallingford CT, Gaussian Inc. **2009**.
33. Lee, C.; Yang, W.; Parr, R. G. *Phys. Rev. B* **1988**, *37*, 785-789 DOI: <https://doi.org/10.1103/PhysRevB.37.785>
34. Becke, A. D. *J. Chem. Phys.* **1993**, *98*, 5648-5652 DOI: <https://doi.org/10.1063/1.464913>
35. Becke, A.D. *J. Chem. Phys.* **1996**, *104*, 1040-1046 DOI: <http://dx.doi.org/10.1063/1.470829>
36. Hay, P. J.; Wadt, W. R. *J. Chem. Phys.* **1985**, *82*, 270-283 DOI: <http://dx.doi.org/10.1063/1.448799>
37. Wadt, W. R.; Hay, P. J. *J. Chem. Phys.* **1985**, *82*, 284-298 DOI: <http://dx.doi.org/10.1063/1.448800>
38. Hay, P. J.;Wadt, W. R. *J. Chem. Phys.* **1985**, *82*, 299-310 DOI: <http://dx.doi.org/10.1063/1.448975>
39. Perrin, D. D. *Stability Constants of Metal-ion Complexes. Part B. Organic Ligand IUPAC Chemical Data Series*, No. 22. Pergamon Press, Oxford, 1979.
40. Arena, G.; Calì, R.; Contino, A.; Musumeci, A.; Musumeci, S.; Purrello, R. *Inorg. Chim. Acta* **1995**, *237*, 187-191 DOI: [https://doi.org/10.1016/0020-1693\(95\)04655-S](https://doi.org/10.1016/0020-1693(95)04655-S)

41. Natsume, T.; Aizawa, S. I.; Hatano, K.; Funahashi, S. *Dalton Trans.* **1994**, 2749-2753 DOI: <https://doi.org/10.1039/DT9940002749>
42. Buzás, N.; Gajda, T.; Nagy, L.; Kuzmann, E.; Vertes, A.; Burger, K. *Inorg. Chim. Acta* **1998**, 274, 167- 176 DOI: [https://doi.org/10.1016/S0020-1693\(97\)06044-1](https://doi.org/10.1016/S0020-1693(97)06044-1)
43. De Stefano, C.; Foti C.; Gianguzza A.; Martino M.; Pellerito L.; Sammartano S. *J. Chem. Eng. Data* **1996**, 41, 511-515 DOI: <https://doi.org/10.1021/je9502915>
44. Foti, C.; Gianguzza, A.; Millero, F.; Sammartano, S. *Aquat. Geochem.* **1999**, 5, 381-398.
45. Rees, D. O. *J. Mol. Biol.* **1980**, 141, 323-326 DOI: [https://doi.org/10.1016/0022-2836\(80\)90184-9](https://doi.org/10.1016/0022-2836(80)90184-9)
46. Rogers, N. K.; Roore, G. R.; Strenberg, M. J. E. *J. Mol. Biol.* **1985**, 182 , 613-616 DOI: [https://doi.org/10.1016/0022-2836\(85\)90248-7](https://doi.org/10.1016/0022-2836(85)90248-7)
47. Sigel, H.; Martin, R. B.; Tribble, R.; Haring, U. K.; Malini-Balakrishnan, R. *Eur. J. Biochem.* **1985**, 152, 187-193 DOI: <https://doi.org/10.1111/j.1432-1033.1985.tb09180.x>
48. Åkerlöf, G.; Short, O. A. *J. Am. Chem. Soc.* **1936**, 58, 1241-1243 DOI: <https://doi.org/10.1021/ja01298a044>
49. Shoukry, M. M.; El-Medani, S. M. *Collect. Czech. Chem. Commun.* **1997**, 62, 1023-1028 DOI: <https://doi.org/10.1135/cccc19971023>
50. Mun, L. C.; Hapipah, M. A.; Shin, S. K. *Appl. Organometal. Chem.* **2012**, 26, 310-319 DOI: <https://doi.org/10.1002/aoc.2862>
51. Al-Flaij, O.; Shehata, M. R.; Mohamed, M. M. A.; Shoukry, M. M. *Monatsh. Chem.* **2001**, 132, 349- 366. DOI: <https://doi.org/10.1007/s007060170>
52. Nath, M.; Singh, H.; Eng, G.; Song, X. *Inorg. Chem. Commun.* **2011**, 14, 1381-1385 DOI: <https://doi.org/10.1016/j.inoche.2011.05.027>