

Diketopiperazines from *Alternaria dauci*

Martha Leyte-Lugo^{1,*}, Pascal Richomme², Luis M. Peña-Rodriguez¹

¹Unidad de Biotecnología, Centro de Investigación Científica de Yucatán, Mérida, Yucatán 97205, Mexico.

²Université d'Angers, UPRES EA921 SONAS, SFR 4207 QUASAV, Angers, France.

***Corresponding author:** Martha Leyte-Lugo, email: : mleyte@correo.xoc.uam.mx

Received March 26th, 2020; Accepted August 14th, 2020.

DOI: <http://dx.doi.org/10.29356/jmcs.v64i4.1228>

Supplementary Information




Fig. S1. ^1H NMR spectrum (600 MHz, CD_3OD) of *cyclo-(pro-val)* (1).




Fig. S2. ^{13}C NMR spectrum (150 MHz, CD_3OD) of *cyclo-(pro-val)* (1).




Fig. S3. ¹H NMR spectrum (400 MHz, CD₃OD) of cyclo-(pro-leu) (2).




Fig. S4. ^{13}C NMR spectrum (100 MHz, CD_3OD) of *cyclo-(pro-leu)* (2).




Fig. S5. ¹H NMR spectrum (600 MHz, CD₃OD) of cyclo-(pro-phe) (3).




Fig. S6. ¹H NMR spectrum (600 MHz, CD₃OD) of cyclo-(val-leu) (4).




Fig. S7. ¹H NMR spectrum (600 MHz, CD₃OD) of cyclo-(val-phe) (5).




Fig. S8. ^1H NMR spectrum (600 MHz, CD_3OD) of cyclo-(leu-phe) (6).




Fig. S9. ¹H NMR spectrum (600 MHz, CD₃OD) of cyclo-(leu-tyr) (7).