

Six-Membered Heterocyclic Boronate Esters. Synthesis and Structural Analysis

Ariana León-Negrete, Raúl Villamil-Ramos, Paola Sánchez-Portillo, Arturo González-Hernández, Victor Barba*

Centro de Investigaciones Químicas-IICBA, Universidad Autónoma del Estado de Morelos, Av. Universidad 1001, C.P. 62209, Cuernavaca, Morelos, México.

*Corresponding author: Victor Barba, email: vbarba@uaem.mx

Received January 11th, 2022; Accepted August 1st, 2022.

DOI: <http://dx.doi.org/10.29356/jmcs.v66i4.1718>

Abstract. Nine heterocyclic zwitterionic boronate esters derived from carbonylphenylboronic acids and amino-diols are described. Compounds were prepared by direct condensation reaction between 3- or 4-formyl/acetylphenylboronic acids with 2-amino-2-methyl-1,3-propanediol (**1a-1d**) or serinol (2-amino-1,3-propanediol) (**1e-1h**). Compound **2e** was obtained by reaction between 4-formylphenylboronic acid and serinol using a solvent mixture methanol/acetone, an aldol condensation reaction was observed. All compounds were characterized by common spectroscopic techniques such as FT-IR, mass spectrometry, and multinuclear ¹H, ¹³C and ¹¹B NMR spectroscopy. ¹¹B NMR spectra showed signals between $\delta = 1.9$ to 7.3 ppm for all compounds, indicating a tetracoordinated environment for the boron atoms in solution. X-ray diffraction analysis showed that boronates are contained in six-membered heterocycles, which have a chair conformation with -OH and -NH₃⁺ substituents in *syn* disposition. The formation of channels in the crystal lattice that are filled with water and supported by hydrogen bonding interactions is noteworthy.

Keywords: Arylboronic acids; zwitterionic species; boronate esters; hydrogen bonds; crystallographic analysis.

Resumen. En el presente trabajo se describen nueve ésteres de boro zwitteriónicos, derivados de ácidos carbonilfenilborónicos. Los compuestos fueron obtenidos mediante reacciones de condensación entre el ácido 3- o 4- formil/acetilfenilborónico con 2-amino-2-metil propanodiol (**1a-1d**) o serinol (**1e-1h**). El compuesto **2e** se sintetizó a través del ácido 4-formilfenilborónico y serinol (2-amino-1,3-propanodiol) utilizando una mezcla de disolventes metanol/acetona, dando lugar a una reacción de condensación aldólica. Los compuestos fueron caracterizados por técnicas espectroscópicas como son FT-IR, espectrometría de masas y espectroscopia multinuclear de RMN ¹H, ¹³C y ¹¹B. El espectro RMN de ¹¹B mostró señales anchas entre $\delta = 1.9$ y 7.3 ppm para todos los compuestos, lo cual indica la presencia de átomos de boro tetracoordinados en solución. El análisis por difracción de rayos-X de monocristal mostró la formación de heterociclos de 6 miembros en conformación *silla*, con una marcada estereoselectividad en donde los grupos -OH y -NH₃⁺ se encuentran en disposición *syn*. En la red cristalina, se observaron canales ocupados por moléculas de agua y soportados por enlaces de hidrógeno.

Palabras clave: Ácidos arilborónicos; especies switteriónicas; ésteres de boro; enlaces de hidrógeno; análisis cristalográfico.

Introduction

It is well known that boronic acids react with 1,2 and 1,3 diols to form 5- or 6-membered cyclic boronic esters, respectively, having the Boron atom in a trigonal planar geometry [1,2]. The boronic acids and diols interactions have received special interest because of their high stability and the directionality offered by the B-O covalent bonds, as well as the reversibility of the process in the presence of an external stimulus such as water molecules, acid or basic media, which promote the hydrolysis of the boronic esters [3-5]. In fact, the characteristics mentioned above, have been exploited in a great variety of applications such as sugar sensing [6-10] or host-guest chemistry [11-13], functional polymers [14], tissue histology [15], NMR shift reagents [16,17], molecular self-assembled materials [18,19], fluorescence imaging carbohydrates [20] and dynamic covalent bonding [21].

A prominent chemical feature of the trivalent boron compounds is the presence of an empty *p*-orbital which is responsible for the Lewis acidity of boron [19], providing the capacity to form coordinative bonds with a series of donor ligands [22,23]. For instance, trisubstituted boron compounds react with N-donor ligands forming N→B coordinative bonds, here the boron atom changes its coordination number to tetracoordinate producing a tetrahedral geometry. The N→B coordinative bond has resulted in a powerful tool for constructing supramolecular architectures such as macrocycles, capsules, and polymers [24-41]. Previously, we described the synthesis of imine boronic esters by multicomponent reactions and demonstrated their ability to sense metal cations conditioned by the presence of a pyridine fragment in the molecular structure [42-44]. In order to get more information about the formation of boronic esters combined with other functional groups such as carbonyl and amine moieties, herein, we report the synthesis and structural analysis of zwitterionic compounds containing boronates which were obtained by esterification reactions.

Experimental

Materials

All reagents and solvents were acquired from commercial suppliers and used without purification.

Instrumentation

The ¹H, ¹³C and ¹¹B NMR spectra were recorded at room temperature using a Varian Gemini 200 or Varian Unit 400 spectrophotometers. TMS (internal, ¹H, $\delta = 0.00$ ppm, ¹³C, $\delta = 0.0$ ppm) and BF₃·OEt₂ (external ¹¹B, $\delta = 0.0$ ppm) were used as standard references. Two-dimensional COSY and HETCOR experiments were carried out for unequivocal signals assignment. The NMR spectra were processed using the MestReNova software [45]. Infrared spectra were recorded on a Nicolet 6700 FT-IR Thermo Scientific spectrophotometer. Mass spectra were obtained with a Jeol JMS 700 equipment. Melting points were determined with a Büchi B-540 digital apparatus.

X-ray Crystallography

X-ray diffraction studies for compounds **1b**, **1d**, **1f** and **2e** were performed on a Bruker-APEX diffractometer with a CCD area detector, using Mo K α -radiation, ($\lambda = 0.71073$) and a graphite monochromator. Frames were collected at $T = 293$ K, by ω/ϕ -rotation ($\Delta/\omega = 0.3^\circ$) at 10 s per frame. The measured intensities were reduced to F^2 . Structure solution, refinement and data output were carried out with the OLEX2 program package [46] using SHELXL-2014 [47] for refinement. Non-hydrogen atoms were refined anisotropically. All hydrogen atoms were placed in geometrically calculated positions using a riding model. Intermolecular distances were analyzed with Mercury [48].

General method for the preparation of heterocyclic boronate esters **1a-1h** and **2e**

Compounds **1a-1h** were synthesized from the equimolecular reaction between the corresponding arylboronic acid derivatives and 2-amino-2-methyl-1,3-propanediol or serinol in methanol (methanol/acetone

for **2e**). For compounds **1a-1h**, the reaction mixtures were stirred for 1 h at room temperature and let stand overnight. The formed white precipitate was collected by filtration, showing low solubility in most common organic solvents. The final products were purified by recrystallization or washing using methanol as solvent. For compound **2e** the reaction was stirred under reflux for 3 h. The product was obtained as yellow crystals after slow evaporation of the solvent.

Compound **1a** was synthesized from 0.05 g (0.33 mmol) of 4-formylphenylboronic acid and 0.035 g (0.33 mmol) of 2-amino-2-methyl-1,3-propanediol in 10 mL of MeOH. The product was obtained as a white powder. Yield 0.070 g (91%); m.p._(decomp) > 300 °C. IR (ATR) ν (cm⁻¹): 3054 (O-H, m), 2864 (N-H, w), 1697 (C=O, m), 1597 (w), 1516 (m), 1381 (w), 1303 (w), 1209 (m), 1172 (m), 1098 (s), 1050 (s), 988 (s), 890 (s), 817 (s), 674 (m). EI-MS m/z (%): 219 ([M-H₂O]⁺, 52), 161 (55), 147 (32), 97 (35), 57 (67). ¹H NMR (200 MHz, D₂O) δ : 9.82 (1H, s, H-C=O), 7.81 (2H, d, J = 7.7 Hz, H-3, H-5), 7.69 (2H, d, J = 7.7 Hz, H-2, H-6), 3.53 (4H, AB, J = 11.0 Hz, H-7, H-9), 1.14 (1H, s, H-10). ¹³C NMR (50 MHz, D₂O) δ : 197.1 (C=O), 133.9 (C-4), 131.8 (C-2, C-6), 129.0 (C-3, C-5), 64.1 (C-7, C-9), 56.4 (C-8), 17.8 (CH₃). ¹¹B NMR (128.4 MHz, D₂O) δ : 2.0 ppm ($h_{1/2}$ = 2133 Hz).

Compound **1b** was synthesized from 0.05 g (0.33 mmol) of 3-formylphenylboronic acid and 0.035 g (0.33 mmol) of 2-amino-2-methyl-1,3-propanediol in 10 mL of MeOH. The product was obtained as a white powder. Yield 0.053 g (69%); m.p. = 195-200 °C. IR (ATR) ν (cm⁻¹): 3092 (O-H, m), 2876 (N-H, w), 1696 (C=O, m), 1572 (w), 1523 (w), 1376 (w), 1311 (w), 1225 (w), 1105 (m), 1058 (s), 988 (s), 898 (s), 829 (s), 792 (s), 699 (s). EI-MS m/z (%): 219 ([M-H₂O]⁺, 100), 187 (59), 147 (39), 114(12), 95(15), 69(23), 57(15). ¹H NMR (200 MHz, DMSO-*d*₆) δ : 9.53 (1H, s, H-C=O), 8.07 (1H, s, H-2), 7.71 (1H, d, J = 7.6 Hz, H-4), 7.62 (1H, d, J = 7.4 Hz, H-6), 7.29 (1H, t, J = 7.6 Hz, H-5), 4.31 (4H, AB, J = 10.7 Hz, H-7, H-9), 1.07 (1H, s, H-10). ¹³C NMR (100 MHz, DMSO-*d*₆) δ : 190.5 (C=O), 136.3 (C-2), 135.7 (C-3), 133.3 (C-6), 130.9 (C-4), 128.3 (C-5), 68.7 (C-7, C-9), 59.5 (C-8), 21.1 (CH₃). ¹¹B NMR (128.4 MHz, D₂O) δ : 5.0 ppm ($h_{1/2}$ = 960 Hz).

Compound **1c** was synthesized from 0.05 g (0.304 mmol) of 4-acetylphenylboronic acid and 0.032 g (0.304 mmol) of 2-amino-2-methyl-1,3-propanediol in 10 mL of MeOH. The product was obtained as a white powder. Yield 0.068 g (89 %); m.p. = 113-119 °C. IR (ATR) ν (cm⁻¹): 3086 (O-H, m), 2873 (N-H, w), 1667 (C=O, m), 1597 (w), 1524 (w), 1393 (w), 1360 (w), 1279 (w), 1205 (w), 1185 (w), 1107 (m), 1054 (s), 993 (m), 862 (s), 825 (s), 747 (m). EI-MS m/z (%): 233 ([M-H₂O]⁺, 100), 191 (25), 186 (29), 161 (53), 114 (14), 57 (17). ¹H NMR (200 MHz, D₂O) δ : 7.86 (2H, d, J = 7.9 Hz, H-3, H-5), 7.63 (2H, d, J = 7.9 Hz, H-2, H-6), 3.54 (4H, AB, J = 12 Hz, H-7, H-9), 2.61 (3H, s, CH₃-), 1.15 (1H, s, H-10). ¹³C NMR (50 MHz, D₂O) δ : 204.9 (C=O), 134.5 (C-4), 131.7 (C-2, C-6), 127.4 (C-3, C-5), 63.8 (C-7, C-9), 56.9 (C-8), 26.7 (CH₃CO), 17.6 (CH₃). ¹¹B NMR (128.4 MHz, D₂O) δ : 3.0 ppm ($h_{1/2}$ = 2560 Hz).

Compound **1d** was synthesized from 0.05 g (0.304 mmol) of 3-acetylphenylboronic acid and 0.032 g (0.304 mmol) of 2-amino-2-methyl-1,3-propanediol in 10 mL of MeOH. The product was obtained as white powder. Yield 0.065 g (85%); m.p. = 75-79 °C. IR (ATR) ν (cm⁻¹): 3091 (O-H, m), 2863 (N-H, w), 1666 (C=O, m), 1585 (w), 1503 (w), 1462 (w), 1430 (w), 1397 (w), 1356 (w), 1258 (m), 1238 (m), 1144 (d), 1111 (s), 1054 (s), 1017 (m), 984 (m), 878 (s), 796 (s), 764 (s), 697 (s). EI-MS m/z (%): 233 ([M-H₂O]⁺, 100), 201 (30), 186 (24), 161 (40), 114 (14), 57 (15). ¹H NMR (200 MHz, DMSO-*d*₆) δ : 8.00 (1H, s, H-2), 7.95 (1H, d, J = 7.0 Hz, H-4), 7.64 (1H, d, J = 7.0 Hz, H-6), 7.52 (1H, t, J = 7.6 Hz, H-5), 3.5 (4H, broad, overlapped with water signal, H-7, H-9), 2.56 (3H, s, CH₃-), 2.42 (1H, s, H-10). ¹³C NMR (50 MHz, DMSO-*d*₆) δ : 198.7 (C=O), 136.5 (C-2), 135.6 (C-3), 133.3 (C-6), 131.4 (C-4), 128.8 (C-5), 71.5 (C-7, C-9), 66.7 (C-8), 26.7 (CH₃CO), 26.5 (CH₃). ¹¹B NMR (128.4 MHz, DMSO-*d*₆) δ : 1.5 ppm ($h_{1/2}$ = 1882 Hz).

Compound **1e** was synthesized from 0.05 g (0.33 mmol) of 4-formylphenylboronic acid and 0.03 g (0.33 mmol) of serinol in 10 mL of MeOH. The product was obtained as a white powder. Yield 0.0505 g (68 %); m.p._{decomp} = 344 °C. IR (ATR) ν (cm⁻¹): 3105 (O-H, m), 2849 (N-H, w), 1671 (C=O, m), 1557 (w), 1476 (m), 1419 (m), 1312 (s), 1244 (s), 1160 (s), 1114 (m), 1064 (m), 1018 (m), 830 (m), 726 (m), 646 (s). EI-MS m/z (%): 205 ([M-H₂O]⁺, 100), 173 (34), 147 (58), 100 (15), 77 (11). ¹H NMR (200 MHz, D₂O) δ : 9.85 (1H, s, H-C=O), 7.77 (4H, m, H-2,6, H-3,5), 3.58-3.86 (5H, m, H-7, H-8, H-9). ¹³C NMR (50 MHz, D₂O) δ : 197.2 (C=O), 134.1 (C-4), 132.2 (C-2, C-6), 129.0 (C-3, C-5), 59.5 (C-7, C-9), 53.8 (C-8). ¹¹B NMR (128.4 MHz, D₂O) δ : 6.6 ppm ($h_{1/2}$ = 358.5 Hz).

Compound **1f** was synthesized from 0.05 g (0.33 mmol) of 3-formylphenylboronic acid and 0.03 g (0.33 mmol) of serinol in 10 mL of MeOH. The product was obtained as a white powder. Yield 0.045 g (60 %);

m.p. = 320-322 °C. IR (ATR) ν (cm⁻¹): 3066 (O-H, m), 2861 (N-H, w), 1686 (C=O, m), 1596 (w), 1543 (w), 1371 (m), 1257 (m), 1199 (m), 1049 (s), 865 (s), 786 (s), 700 (s), 647 (s). EI-MS m/z (%): 205 ([M-H₂O]⁺, 32), 175 (100), 119 (22), 91(22), 57(33). ¹H NMR (200 MHz, DMSO-*d*₆) δ : 9.90 (1H, s, H-C=O), 8.11 (1H, s, H-2), 7.93 (1H, d, *J* = 7.5 Hz, H-4), 7.80 (1H, d, *J* = 7.5 Hz, H-6), 7.50 (1H, t, *J* = 7.45 Hz, H-5), 3.54-3.78 (5H, m, H-7, H-8, H-9). ¹³C NMR (50 MHz, DMSO-*d*₆) δ : 197.9 (C=O), 139.3 (C-2), 134.8 (C-3), 134.2 (C-6), 128.4 (C-4), 128.2 (C-5), 60.4 (C-7, C-9), 53.5 (C-8). ¹¹B NMR (128.4 MHz, D₂O) δ : 7.3 ppm (*h*_{1/2} = 256.8 Hz).


Compound **1g** was synthesized from 0.05 g (0.304 mmol) of 4-acetylphenylboronic acid and 0.027 g (0.304 mmol) of serinol in 10 mL of MeOH. The product was obtained as a white powder. Yield 0.049 g (69 %); m.p. = 109-111 °C. IR (ATR) ν (cm⁻¹): 3089 (O-H, m), 2871 (N-H, w), 1674 (C=O, m), 1598 (w), 1541 (w), 1390 (w), 1372 (w), 1353 (w), 1278 (w), 1254 (w), 1186 (w), 1125 (w), 1050 (m), 957 (w), 861 (s), 827 (s), 800 (m), 747 (m), 629 (m). EI-MS m/z (%): 219 ([M-H₂O]⁺, 100), 204 (25), 187 (10), 172 (15), 161 (46), 120 (11), 100 (10). ¹H NMR (200 MHz, D₂O) δ : 7.88 (2H, d, *J* = 7.50 Hz, H-3, H-5), 7.68 (2H, d, *J* = 7.5 Hz, H-2, H-6), 3.62-4.02 (5H, m, H-7, H-8, H-9), 2.63 (3H, s, CH₃). ¹³C NMR (50 MHz, D₂O) δ : 204.9 (C=O), 134.9 (C-4), 132.0 (C-2, C-6), 127.4 (C-3, C-5), 60.0 (C-7, C-9), 53.8 (C-8), 26.0 (CH₃CO). ¹¹B NMR (128.4 MHz, D₂O) δ : 2.0 ppm (*h*_{1/2} = 513.6 Hz).

Compound **1h** was synthesized from 0.05 g (0.304 mmol) of 3-acetylphenylboronic acid and 0.027 g (0.304 mmol) of serinol in 10 mL of MeOH. The product was obtained as a white powder. Yield 0.049 g (68 %); m.p. = 120-121 °C. IR (ATR) ν (cm⁻¹): 3073 (O-H, m), 2875 (N-H, w), 1675 (C=O, m), 1510 (m), 1428 (w), 1392 (w), 1361 (w), 1268 (m), 1248 (m), 1148 (m), 1110 (m), 1043 (m), 989 (m), 873 (s), 821 (s), 737 (s), 622 (s). EI-MS m/z (%): 219 ([M-H₂O]⁺, 100), 204 (63), 187 (30), 172 (48), 161 (94), 146 (17), 131 (14), 120 (31), 103 (24), 100 (34). ¹H NMR (200 MHz, D₂O) δ : 8.18 (1H, s, H-2), 7.84 (2H, m, H-4, H-6), 7.44 (1H, t, *J* = 7.7 Hz, H-5), 3.65-3.93 (5H, m, H-7, H-8, H-9), 2.67 (3H, s, CH₃). ¹³C NMR (50 MHz, D₂O) δ : 205.5 (C=O), 137.8 (C-2), 135.4 (C-3), 131.9 (C-6), 127.7 (C-4), 126.8 (C-5), 60.2 (C-7, C-9), 53.6 (C-8), 25.6 (CH₃CO). ¹¹B NMR (128.4 MHz, DMSO-*d*₆) δ : 1.8 ppm (*h*_{1/2} = 924.5 Hz).


Compound **2e** was synthesized from 0.184 g (1.23 mmol) of 4-formylphenylboronic acid and 0.112 g (1.23 mmol) of serinol in 30 mL of MeOH/Acetone (1:1 V/V) solvent mixture. Yield 0.104 g (32 %); m.p. = 103-105 °C. IR (ATR) ν (cm⁻¹): 3103 (O-H, m), 2869 (N-H, w), 1682 (C=O, m), 1621 (C=C, m) 1556 (w), 1413 (w), 1318 (m), 1253 (s), 1151 (m), 1125 (m), 1047 (s), 1015 (s), 970 (s), 896 (s), 865 (s), 810 (s), 737 (s), 643 (s), 634 (s). EI-MS m/z (%): 245 ([M-H₂O]⁺, 45), 230 (84), 187 (100), 145 (35), 129 (49), 77 (38), 60 (53). ¹H NMR (200 MHz, DMSO-*d*₆) δ : 7.56-7.86 (5H, m, H-2,6, H-3,5, H-10), 6.83 (1H, d, ³*J*_{trans} = 16.5 Hz, H-11), 3.56-3.97 (5H, broad, overlapped with water signal, H-7, H-8, H-9), 2.32 (3H, s, H-13). ¹³C NMR (50 MHz, DMSO-*d*₆) δ : 198.2 (C=O), 144.7 (COCH), 134.6 (C-4), 133.4 (C-2, C-6), 132.3 (C-3, C-5), 127.2 (CH-Ar), 62.8 (C-7, C-9), 54.6 (C-8), 27.4 (CH₃CO). ¹¹B NMR (128.4 MHz, D₂O) δ : 1.1 ppm (*h*_{1/2} = 61.14 Hz).

Results and discussion

Zwitterionic boronate esters were prepared in a reaction of carbonylphenylboronic acid derivatives (4-formylphenylboronic acid; **1a**, **1e**), (3-formylphenylboronic acid; **1b**, **1f**), (4-acetylphenylboronic acid; **1c**, **1g**), (3-acetylphenylboronic acid; **1d**, **1h**) with 2-amino-2-methyl-1,3-propanediol (**1a-1d**) or serinol (**1e-1h**), in a 1:1 equimolar ratio using methanol as solvent (Scheme 1). The reaction mixtures were stirred at room temperature for 1 hour and upon standing over one night, a white precipitate appeared. The products were recovered by filtration and purified by crystallization from methanol resulting in moderate yields (~60 %). When the reaction of 4-formylphenylboronic acid with serinol in a 1:1 equimolar ratio was carried out using acetone/methanol as a solvent mixture (Scheme 2), compound **2e** was obtained as a yellow pale solid in low yield (32 %). Besides the boronate ester formation, an α,β -unsaturated ketone was obtained as a product of the aldol condensation reaction involving the aldehyde group and the acetone solvent.



Scheme 1. Synthesis of the zwitterionic boronate esters **1a-1h**.



Scheme 2. Synthesis of the zwitterionic boronate ester **2e** showing the aldol condensation reaction.

In all cases, the boronic acid reacts with the amino-diol and forms the boronic ester, and while the carbonyl groups were supposed to react with the amine group forming an imine bond, the boronic ester reacted with a water molecule forming a zwitterionic derivative. Thus, the amine group (NH_2) is protonated yielding an ammonium group (NH_3^+) and the hydroxyl group ($-\text{OH}$) is attached to the boron atom forming a boronate anion.

Spectroscopic analysis

All compounds **1a-1h** and **2e** were characterized by IR spectroscopy, mass spectrometry and ^1H , ^{13}C and ^{11}B NMR spectroscopy. Furthermore, the structures of four zwitterionic boronate esters (**1b**, **1d**, **1f** and **2e**) were established by single crystal X-ray diffraction. The first approximation to characterize the compounds was carried out by IR spectroscopy. In the IR spectra, all compounds **1a-1h** and **2e** showed broad absorption bands in the range from 3054 to 3105 cm^{-1} due to the presence of hydroxyl groups $\nu(\text{OH})$, while the bands observed between 2849 to 2876 cm^{-1} correspond to the ammonium groups. Moreover, the band observed at 1666 - 1697 cm^{-1} was assigned to the stretching vibrations of the carbonyl group $\nu(\text{C}=\text{O})$. Additionally, a new absorption band for compounds **1a-1h** and **2e** appear at 1043 - 1064 cm^{-1} which is characteristic of the stretching vibration of the $\nu(\text{B}-\text{O})$ group. The formation of six-membered heterocyclic boronates **1a-1h** was confirmed by EI-MS, where the spectra showed peaks at $m/z = 219$ (**1a,1b**), 233 (**1c,1d**), 205 (**1e,1f**), 219 (**1g, 1h**) and 245 (**2e**) corresponding to the molecular ion $[\text{M}]^+$ with the loss of a water molecule. In most of the cases, this peak corresponds to the base peak indicating good stability of the compounds.

The characterization of compounds **1a-1h** and **2e** was complemented by NMR analyses. All compounds showed low solubility in common organic solvents. NMR spectra were recorded using $\text{DMSO}-d_6$ (**1b**, **1d** and **2e**) and D_2O (**1a**, **1c**, **1e**, and **1h**). The ^1H NMR spectra of the formylboronic acid derivatives show

signals at high frequency, between $\delta = 9.53$ - 9.90 ppm assigned to the hydrogen atom of the aldehyde group. The signal for methylene hydrogens (CH_2 , H-7, H-9) for 2-amino-2-methyl-1,3-propanediol derivatives (**1a-1d**) was observed as an *AB* system at $\delta = 3.50$ - 4.31 ppm, having coupling constants from 10.7 to 12.0 Hz. While for compounds **1e-1h** and **2e**, the signals corresponding to methylene hydrogens are observed as multiples and shifted between $\delta = 3.54$ - 4.02 ppm. The ^{13}C NMR also showed the corresponding signals for the carbonyl group which appears in the range of $\delta = 198.7$ - 205.5 ppm for acetyl derivatives, while for aldehyde derivatives the signals were observed at a lower frequency (198.2 - 190.5 ppm). The carbon signals for aliphatic groups of the heterocyclic ring were observed between $\delta = 53.5$ - 71.5 ppm. Finally, the ^{11}B NMR spectra of all the compounds display a single broad signal in the range of $\delta = 1.9$ - 7.3 ppm, as expected for tetracoordinated boron derivatives [35-39].

Suitable crystals for X-ray diffraction analysis were obtained for compounds **1b**, **1d**, **1f** and **2e** (Fig. 1) by recrystallization process from methanol solutions, crystallographic data are described at Table 1. Compounds **1d**, **1f** and **2e** crystallized in monoclinic system, while **1b** crystallized in a triclinic system. The asymmetric unit shows water molecules surrounding the boronate molecules, two for **1b**, **1f**, **2e**, and four and a half for **1d**.




Fig. 1. Molecular structures for compounds **1b**, **1d**, **1f** and **2e** (50 % ellipsoid probability).

Table 1. Selected crystallographic data for compounds **1b**, **1d**, **1f** and **2e**.

	1b	1d	1f	2e
Empirical formula	$\text{C}_{11}\text{H}_{19}\text{BNO}_6$	$\text{C}_{12}\text{H}_{26}\text{BNO}_{8.5}$	$\text{C}_{10}\text{H}_{17}\text{BNO}_6$	$\text{C}_{13}\text{H}_{22}\text{BNO}_6$
Fw	272.08	331.15	258.05	299.12
Cryst size [mm^3]	0.15 x 0.24 x 0.32	0.22 x 0.27 x 0.45	0.23 x 0.36 x 0.40	0.22 x 0.36 x 0.44
Cryst syst	Triclinic	Monoclinic	Monoclinic	Monoclinic
Space group	P1	C2/c	P2 ₁ /c	P2 ₁ /c
Unit cell dimensions				
a [\AA]	6.5199(15)	31.44(2)	15.319(4)	17.490(6)
b [\AA]	7.7679(18)	9.062(6)	12.176(4)	12.586(4)
c [\AA]	14.562(3)	12.104(8)	6.908(2)	6.842(2)
α [$^\circ$]	103.886(4)	90	90	90
β [$^\circ$]	90.806(4)	94.275(10)	96.023(5)	94.147(6)
γ [$^\circ$]	103.850(4)	90	90	90
V [\AA^3]	693.2(3)	3439(4)	1281.3(6)	1502.1(9)
Z	2	8	4	4
ρ_{calcd} [Mg/m^3]	1.304	1.279	1.338	1.323

$\mu[\text{mm}^{-1}]$	0.104	0.106	0.108	0.102
Collected reflns	2912	8993	11141	12307
Independent reflns (R_{int})	1892 (0.0275)	2471 (0.0347)	2001 (0.0834)	2220 (0.1213)
Number of param	187	240	190	209
$R_1 [I > 2\sigma(I)]$	0.0751	0.0462	0.0497	0.0612
wR2 (all data)	0.1932	0.1257	0.1380	0.1557
GOF	1.083	1.119	1.074	0.900

Crystals for compounds **1a**, **1c**, **1e** and **1g** were also obtained (Fig. 2), however most of them include several disorder water molecules at the unit cell, for that reason, refinement of the structures was inadequate or included high R index values (Table S1). Nevertheless, the X-ray diffraction analysis allowed the observation of the whole molecular structures (only connective models).





Fig. 2. Connective models for compounds **1a**, **1c**, **1e** and **1g**.

The X-ray diffraction analysis shows a chair conformation for the six-membered heterocyclic ring for all eight compounds (Fig. 1). In fact, the presence of the $-\text{OH}$ and $-\text{NH}_3^+$ groups at the six-membered ring gives the possible *anti* and *syn* isomers, resulting in four possible conformations (I-IV, Scheme 3). In this work, it was noticed, that the aryl and ammonium groups occupied the axial positions, whereas the methyl/proton and hydroxyl groups are in the equatorial positions. The above observation shows the preference for the *syn* stereoisomer considering the relationship between the hydroxyl and ammonium groups (isomer III at Scheme 3).

The endocyclic O-B bond distances are very similar (1.481(4) to 1.498(3) Å), being longer than the observed for the exocyclic B-O bond distances [1.453(3) -1.465(4) Å], values are related to similar reported boronate compounds [35-37, 49-51]. The angles observed around the boron atom are closer to the ideal value in a tetrahedral geometry (109.5°) having values of 104.1(4) to 113.9(4) °. In fact, the tetrahedral character (THC) for the boron atoms showed values of 80.5, 87.0 and 86.7 %, respectively [38,52], indicating the *pseudo* tetrahedral geometry (Table 2).



Scheme 3. Possible *syn*- and *anti*-conformations for boronates **1a-1h**.

Table 2. Selected bond lengths (Å), angles (°) and THC for compounds **1b**, **1d**, **1f** and **2e**.

Bonds (Å)				
Compound	1b	1d	1f	2e
B(1)-O(1)	1.495(3)	1.498(3)	1.481(4)	1.493(4)
B(1)-O(2)	1.493(6)	1.497(3)	1.494(4)	1.497(4)
B(1)-O(3)	1.460(6)	1.453(3)	1.465(4)	1.45(4)
B(1)-C _{Ar}	1.628(7)	1.623(4)	1.626(4)	1.623(5)
Angles (°)				
O(3)-B(1)-O(2)	108.8(4)	106.8(2)	107.7(2)	1091(3)
O(3)-B(1)-O(1)	104.1(4)	107.06(19)	108.6(2)	1076(3)
O(2)-B(1)-O(1)	106.6(4)	108.56(19)	108.1(2)	1070(3)
O(3)-B(1)-C _{Ar}	112.2(4)	110.5(2)	108.0(2)	1089(3)
O(2)-B(1)-C _{Ar}	110.9(4)	112.5(2)	112.3(2)	1104(3)
O(1)-B(1)-C _{Ar}	113.9(4)	111.1(2)	113.0(2)	113.8(3)
THC(%)	80.5	87.0	86.7	88.2

As mentioned above, compounds **1a-1g** crystallized with water molecules, which interact by hydrogen bonds with the boronates through the -OH and NH_3^+ groups. The supramolecular analysis for compounds **1b**, **1d**, and **1f** in solid-state reveals the presence of several hydrogen bonds that result in the formation of 1D polymers. For instance, compound **1d** is surrounded by seven water molecules *via* $\text{N-H}\cdots\text{O}$ and $\text{O-H}\cdots\text{O}$ interactions as depicted in Fig. 3(a). Moreover, it was observed the presence of six water molecules located between the donor and acceptor groups (-NH_3^+ , -OH), closing the interaction face to face of two boronate molecules as depicted in Fig. 3(b). The above observation favors the *syn* orientation of the OH and NH_3^+ groups as described previously at Scheme 3 (III). For compound **1d**, the distances for the $\text{N-H}\cdots\text{O}$ donor-acceptor interactions have values of 1.962, 1.973 and 2.214 Å, whereas the $\text{O-H}\cdots\text{O}$ interaction has a distance of 2.014 Å. Fig. 3(c) and 3(d) show that the hydrogen bonds between the ammonium or hydroxyl groups with water molecules, support the formation of 1D polymer chains (Fig. S1), and give place to the generation of water channels along *b*-axis. Similar water chains were observed for compounds **1b** and **1f** (Figures S2 and S3) and analogous channels have been described for boron compounds [53] and hydrophilic organic molecules [54-56].




Fig. 3. (a) Seven water molecules have direct interaction with the boronate **1d** via N-H...O and O-H...O hydrogen bonds. (b) Six water molecules connect face to face two boronate units. (c) Crystal packing for compound **1d** along *b*-axis showing the polymeric chains of boronate and water molecules. (d) Crystal packing for compound **1d** along *a*-axis.

Suitable crystals for the X-ray analysis were also obtained for compound **2e** after filtration and slow evaporation of the mother solution. As mentioned above, complex **2e** resulted from the aldol condensation reaction between the formyl group of **1e** and the acetone solvent, giving rise to the formation of the respective carbon-carbon double bond in an *E* configuration. In fact, structural parameters related to the six-membered heterocycle conformation remain without significant changes, and bond distances and angles around the boron atom in complex **2e** are comparable to its partners **1b**, **1d**, and **1f**. The endocyclic B-O bond distances are slightly longer (1.493(4) - 1.497(4) Å) compared with the exocyclic B-O distance (1.455(4) Å) of the hydroxyl group. The angles around the boron atom are between 107.0 to 113.8° with a THC value of 88.2 %, confirming the *pseudo* tetrahedral geometry. The -NH₃⁺ and -OH groups preserve the *syn* conformation in the six-membered ring.

The supramolecular analysis for **2e** reveals that each boronate molecule interacts with four water molecules via N-H...O and O-H...O hydrogen bonds (Fig. 4(a)). Furthermore, four water molecules connect two boronate units by hydrogen bonds (Fig. 4(b)). The distances for the N-H...O donor-acceptor interactions have values of 1.887, 1.986 and 2.090 Å, whereas the O-H...O interaction has a distance of 1.978 Å. It was noticed that the presence of α,β -unsaturated substituent does not affect the molecular packing, which looks like the observed for **1d**. Once more, the observable *1D* polymer chain for the boronate units results from the presence of water molecules connected via N-H...O and O-H...O hydrogen bonds (Figures 4(c) and S4).




Fig. 4. (a) Interaction of compound **2e** interacting with four water molecules via N-H...O and O-H...O hydrogen bonds. (b) Two boronate units connected by four water molecules. (c) Packing in compound **2e** along *b*-axis showing the polymeric chains including channels water formation.

Conclusion

Nine heterocyclic boron compounds were prepared by the condensation reaction of 2-amino-1,3-propanediol derivatives with carbonylphenylboronic acids at room temperature. Spectroscopic characterization indicated the formation of zwitterionic boronates, including the ammonium (NH₃⁺) and hydroxyl (OH) groups. The zwitterionic character decrease the solubility of compounds in common organic solvents, so, the possible reaction between the amino and carbonyl groups is inhibit, and a further self-assembly reaction was not achieved. Molecular structures obtained by X-ray diffraction analysis showed a chair conformation for the six-membered ring having the -NH₃⁺ and -OH groups in a *syn* disposition. The presence of water molecules in the crystal network favored the formation of *1D* polymeric chains supported by N-H...O and O-H...O hydrogen bonds. Using as solvent acetone, the aldol condensation was observed, while the boronate unit remains without modification; this reactivity opens the possibility to construct boronic acids with extending unsaturated moieties, which could be useful as sensors/indicators of anions, cations or neutral molecules.

Acknowledgments

The authors thank to Consejo Nacional de Ciencia y Tecnología (CONACyT) for financial support (project number A1-S-33602). Authors also acknowledge to CIQ-LANEM for structural analysis.

References

1. Severin, K. *Dalton Trans.* **2009**, 5254-5264.
2. Höpfl, H. *Struct. Bonding.* **2002**, 103, 1-56.
3. Wang, W.; Gao, X.; Wang, B. *Curr. Org. Chem.* **2002**, 6, 14, 1285-1317.
4. Springsteen, G.; Wang, B. *Tetrahedron.* **2002**, 58, 26, 5291-5300.
5. Marinez-Aguirre, M. A.; Villamil-Ramos, R.; Guerrero-Alvarez, J. A.; Yatsimirsky A. K. *J. Org. Chem.* **2013**, 78, 10, 4674-4684.

6. Hall, D. G. *Boronic Acids: Preparation and Applications in Organic Synthesis Medicine and Materials*, 2nd ed, Wiley-VCH, Weinheim, **2011**.
7. James, T. D; Philips, M. D; Shinkai, S. *Boronic acids in Saccharide Recognition* Recognition, The Royal Society of Chemistry, **2006**.
8. James, T.D., Shinkai, S. Artificial Receptors as Chemosensors for Carbohydrates. In: Penadés, S. (eds) *Host-Guest Chemistry. Topics in Current Chemistry*, Vol. 218, Ed., Springer, Berlin, Heidelberg, **2002**.
9. James, T. D; Sandanayake, K. R. A; Shinkai, S. *Angew. Chem. Int. Ed. Engl.* **1996**, 35, 1910-1922.
10. Jin, S; Cheng, Y; Reid, S; Li, M; Wang, B. *Med. Res. Rev.* **2010**, 30, 171-257.
11. Rowan, S. J; Cantrill, S. J; Cousins, G. R; Sanders, J. K; Stoddart, J. F. *Angew. Chem. Int. Ed.* **2002**, 41, 898-952.
12. Corbett, P.T.; Leclaire, J.; Vial, L.; West, K.R.; Wietor, J.L.; Sanders, J. K. M.; Otto, S. *Chem. Rev.* **2006**, 106, 3652-3711.
13. Wilson, A; Gasparini, G; Matile, S. *Chem. Soc. Rev.* **2014**, 43, 1948-1962.
14. Cambre, J. N; Sumerlin, B. S. *Polymer.* **2011**, 52, 4631-4643.
15. Dai, C; Cazares, L. H; Wang, L; Chu, Y; Wang, S. L; Troyer, D. A; Semmes, O. J; Drake, R; Wang, B. *Chem. Commun.* **2011**, 47, 10338-10340.
16. Kelly, A. M; Pérez-Fuertes, Y; Arimori, S; Bull, S. D; James, T. D. *Org. Lett.* **2006**, 8, 1971-1974.
17. Pérez-Fuertes, Y; Kelly, A. M; Johnson, A. L; Arimori, S; Bull, S. D; James, T. D. *Org. Lett.* **2006**, 8, 609-612.
18. Fujita, N; Shinkai, S; James, T. D. *Chem. Asian J.* **2008**, 3, 1076-1091.
19. Nishiyabu, T; Kubo, Y; James, T. D; Fossey, J. S. *Chem. Commun.* **2011**, 47, 1124-1150.
20. Sun, X; Zhai, W; Fossey, J. S; James, T. D. *Chem. Commun.* **2016**, 52, 3456-3469.
21. Bull, S. D; Davidson, M. G; van den Elsen, J. M. H; Fossey, J. S; Jenkins, A. T. A; Jiang, Y.-B; Kubo, Y; Marken, F; Sakurai, K; Zhao, J; James, T. D. *Acc. Chem. Res.* **2013**, 46, 312-326.
22. Carboni, B; Monnier, L. *Tetrahedron.* **1999**, 55, 1197-1248.
23. Vargas-Díaz, G; Höpfl, H. *J. Organomet. Chem.* **2009**, 694, 3660-3666.
24. Farfán, N; Höpfl, H; Barba, V; Ochoa, M. E; Santillan, R; Gómez, E; Gutiérrez, A. *J. Organomet. Chem.* **1999**, 581, 70-81.
25. Höpfl, H; Sánchez, M; Farfán, N; Barba, V. *Can. J. Chem.* **1998**, 76, 1352-1360.
26. Kubo, Y; Nishiyabu, R; James, T.D. *Chem. Commun.* **2015**, 51, 2005-2020
27. Salazar-Mendoza, D; Cruz-Huerta, J; Höpfl, H; Ahuactzi, I. F; Sánchez, M. *Cryst. Growth Des.* **2013**, 13, 2441-2454.
28. Sheepwash, E; Zhou, K; Scopelliti, R; Severin, K. *Eur. J. Inorg. Chem.* **2013**, 2013, 2558-2563.
29. Barba, V; Betanzos, I. *J. Organomet. Chem.* **2007**, 692, 4903-4908.
30. Barba, V; Villamil, R; Luna, R; Godoy-Alcántar, C; Höpfl, H; Beltran, H. I; Zamudio-Rivera, L. S; Santillan, R; Farfán, N. *Inorg. Chem.* **2006**, 45, 2553-2561.
31. Sheepwash, E; Luisier, N; Krause, M. R; Noé, S; Kubik, S; Severin, K. *Chem. Commun.* **2012**, 48, 7808-7810.
32. Sheepwash, E; Kraml, V; Scopelliti, R; Sereda, O; Neels, A; Severin, K. *Angew. Chem. Int. Ed.* **2011**, 50, 3034-3037.
33. Christinat, N; Scopelliti, R; Severin, K. *Angew. Chem. Int. Ed.* **2008**, 47, 1848-1852.
34. Nishimura, N; Yoza, K; Kobayashi, K. *J. Am. Chem. Soc.* **2010**, 132, 777-790.
35. Barba, V; Höpfl, H; Farfán, N; Santillan, R; Beltran, H. I; Zamudio-Rivera, L. S. *Chem. Commun.* **2004**, 2834-2835.
36. Barba, V; Ramos, P; Jiménez, D; Rivera, A; Meneses, A. *Inorg. Chim. Acta.* **2013**, 401, 30-37.
37. Celis, N. A; Godoy-Alcántar, C; Guerrero-Álvarez, J; Barba, V. *Eur. J. Inorg. Chem.* **2014**, 1477-1484.
38. Gómez-Jaimes, G; Barba, V. *J. Mol. Struct.* **2014**, 1075, 594-598.
39. González-Hernández, A; Serrano-Melgar, G; Villamil-Ramos, R; Barba, V. *Heteroat. Chem.* **2017**, 28, e21377.

40. Herrera-España, A.D.; Campillo-Alvarado, G.; Román-Bravo, P.; Herrera-Ruiz, D.; Höpfl, H.; Morales-Rojas, H. *Cryst. Growth Des.* **2015**, *15*, 4, 1572-1576.
41. Campillo-Alvarado, G.; Vargas-Olvera, E. C.; Höpfl, H.; Herrera-España, A. D.; Sánchez-Guadarrama, O.; Morales-Rojas, H.; MacGillivray, L. R.; Rodríguez-Molina, B.; Fáfán N.; *Cryst. Growth Des.* **2018**, *18*, 5, 2726-2743.
42. Sanchez-Portillo, P; Arenaza-Corona, A; Hernandez-Ahuactzi, I. F; Barba, V. *J. Mol. Struct.* **2017**, *1134*, 435-443.
43. Sanchez-Portillo, P; Barba, V. *ChemistrySelect.* **2017**, *2*, 11265-11272.
44. Sánchez-Portillo, P; Hernández-Sirio, A; Godoy-Alcantar, C; Lacroix, P. G; Agarwal, V; Santillán, R; Barba, V. *Dyes Pigment.* **2021**, *186*, 108991.
45. MestReNova, N° de versión 12.0.0, **2017**, Windows. Mexico: Mestrelab Research.
46. Dolomanov, V; Bourhis, L. J; Gildea, R. J; Howard, J. A. K; Puschmann, H. *J. Appl. Cryst.* **2009**, *42*, 339-341.
47. Sheldrick, G. M. *Acta Cryst C.* **2015**, *71*, 3-8.
48. Edgington, P. R; McCabe, P; Macrae, C. F; Pidock, E; Shields, G. P; Taylor, R; Towler, M; Van De Streek, J. *J. Appl. Crystallogr.* **2006**, *39*, 453-457.
49. Höpfl, H; Farfán, N. *J. Organomet. Chem.* **1997**, *547*, 71-77.
50. Barba, V; Vargas, G; Gómez, E; Farfán, N. *Inorg. Chim. Acta.* **2000**, *311*, 133-137.
51. Rivera, J. M; Rincón, S; Farfán, N; Santillan, R. *J. Organomet. Chem.* **2011**, *696*, 2420-2428.
52. Höpfl, H. *J. Organomet. Chem.* **1999**, *581*, 129-149.
53. Rodríguez-Cuamatzi, P; Vargas-Díaz, G; Höpfl, H. *Angew. Chem. Int. Ed.* **2004**, *43*, 3041-3044.
54. Wang, J; Zheng, L.L; Li, C. J; Zheng, Y. Z; Tong, M. L. *Cryst. Growth Des.* **2006**, *6*, 357-359.
55. Wang, Y. T; Tang, G. M; Liu, Z. M; Yi, X. H. *Cryst. Growth Des.* **2007**, *7*, 2272-2275.
56. Natarajan, R; Charmant, J. P. H; Orpen, A. G; Davis, A. P. *Angew. Chem. Int. Ed.* **2010**, *49*, 5125-5129.